



Review

Nutrient constraints on terrestrial carbon fixation: The role of nitrogen[☆]Devrim Coskun, Dev T. Britto, Herbert J. Kronzucker^{*}

Department of Biological Sciences and the Canadian Centre for World Hunger Research (CCWHR), University of Toronto, Canada

ARTICLE INFO

Article history:

Received 4 March 2016

Received in revised form 26 May 2016

Accepted 30 May 2016

Available online 11 June 2016

Keywords:

Carbon dioxide (CO₂)

Nitrogen (N)

Nitrate (NO₃⁻)Ammonium (NH₄⁺)

Photosynthesis

Respiration

Metabolism

Climate change

ABSTRACT

Carbon dioxide (CO₂) concentrations in the earth's atmosphere are projected to rise from current levels near 400 ppm to over 700 ppm by the end of the 21st century. Projections over this time frame must take into account the increases in total net primary production (NPP) expected from terrestrial plants, which result from elevated CO₂ (eCO₂) and have the potential to mitigate the impact of anthropogenic CO₂ emissions. However, a growing body of evidence indicates that limitations in soil nutrients, particularly nitrogen (N), the soil nutrient most limiting to plant growth, may greatly constrain future carbon fixation. Here, we review recent studies about the relationships between soil N supply, plant N nutrition, and carbon fixation in higher plants under eCO₂, highlighting key discoveries made in the field, particularly from free-air CO₂ enrichment (FACE) technology, and relate these findings to physiological and ecological mechanisms.

© 2016 Elsevier GmbH. All rights reserved.

Contents

1. Introduction.....	96
2. N constraints on plant-growth responses to eCO ₂	96
2.1. Methodological note.....	96
2.2. Grasslands.....	97
2.3. Forests.....	99
2.4. Agriculture.....	100
3. Plant and soil N uptake and accumulation under eCO ₂	101
3.1. Tissue N content.....	101
3.2. The effects of contrasting N sources.....	101
3.3. Plant N acquisition.....	102
3.4. Progressive N limitation (PNL) and N deposition.....	102
3.5. Biological N ₂ fixation (BNF).....	104
4. Conclusion.....	104
Acknowledgements.....	104
References.....	105

Abbreviations: A', daily integrated CO₂ uptake; A_{net}, net photosynthesis; A_{sat}, light-saturated CO₂ uptake; BNF, biological N₂ fixation; C_a, atmospheric CO₂ concentrations; CUE, ecosystem carbon-use efficiency; eCO₂, elevated atmospheric CO₂; FACE, free-air CO₂ enrichment; GPP, gross primary production; g_s, stomatal conductance; MSX, methionine sulfoximine; NEP, net ecosystem production; N_m, concentration of N in plant dry mass; NPP, net primary production; N_r, reactive nitrogen; OTC, open-top chambers; PEPC, phosphoenolpyruvate carboxylase; PNL, progressive nitrogen limitation; PNUE, photosynthetic N-use efficiency; SLA, specific leaf area; SOM, soil organic matter.

[☆] This article is part of a special section entitled "Plants facing Changing Climate", published in the Journal of Plant Physiology 203, 2016.

^{*} Corresponding author.

E-mail address: herbert.kronzucker@utoronto.ca (H.J. Kronzucker).

1. Introduction

Atmospheric CO₂ concentrations (C_a) have increased by nearly 50% since pre-industrial times, to values currently approaching 400 ppm. Socio-biogeochemical models have predicted that C_a may reach 550 ppm by the middle of the century, and may well surpass 700 ppm by 2100 (Pachauri et al., 2014; Le Quéré et al., 2015). Such levels have not been known to occur on Earth since the early Miocene, some 24 million years ago (Pagani et al., 2009). Closely associated with these forecasts are predictions of disastrous changes in the Earth's climate (e.g. Yin, 2013; Cai et al., 2014), even if all anthropogenic emissions were to halt today (Frölicher et al., 2014). Accordingly, much recent attention has been focused on the consumption and storage of CO₂ by terrestrial biomes, where as much as a third of all anthropogenic CO₂ emissions (currently ~10 Pg C yr⁻¹) is captured (Reay et al., 2008; Andres et al., 2012; Le Quéré et al., 2015).

It is well established that elevated atmospheric CO₂ (eCO₂) can stimulate net primary production (NPP) in photosynthetic organisms (Harley et al., 1992; Norby et al., 2005; McCarthy et al., 2010; Franks et al., 2013; see also Fig. 1). Indeed, crop cultivation in greenhouses routinely involves the enrichment of greenhouse air with CO₂, which is applied at levels as high as 1000 ppm, and results in growth and yield increases of as much as 30% (Becker and Kläring, 2016). In field experiments, eCO₂ (at ~550 ppm) has been shown to generally increase carbon gain and biomass increases in a wide range of C₃ plant systems, by amounts that vary from study to study. For instance, in a meta-analysis of FACE (Free-Air CO₂ Enrichment) studies, Leakey et al. (2009) found biomass increases of about 19–46%, compared to growth under present-day CO₂ conditions. Lee et al. (2011), examining 13 grassland species, found a more modest increase of about 10%. By contrast, the photosynthetic apparatus of C₄ plants is already saturated under current CO₂ concentrations, and therefore elevated CO₂ does not have a great impact on the growth of this functional group (Lee et al., 2011; Kant et al., 2012), which includes important agricultural crops such as corn and sorghum. In addition to increases in plant biomass due to a 'CO₂ fertilization effect', eCO₂ has been linked to stimulated photosynthetic output, reduced stomatal conductance and transpiration, and increases in the efficiencies of water, light, and nitrogen (N) use (Curtis and Wang, 1998; Drake et al., 1997; Leakey et al., 2009).

However, constraints imposed on eCO₂-enhanced carbon fixation and plant productivity due to soil nutrient limitations, particularly that of N, have also long been observed (Evans, 1989; Vitousek and Howarth, 1991; Drake et al., 1997; LeBauer and Treseder, 2008). Higher rates of growth lead to increased demand for nutrients, especially N, which represents the most frequently growth-limiting soil nutrient in terrestrial ecosystems (cf. Sardans and Peñuelas, 2015). Moreover, although initially elevated rates of NPP under eCO₂ become downregulated in the longer term, involving the process of photosynthetic acclimation (Sage, 1994; Stitt and Krapp, 1999; see also Section 3; Fig. 2), they decrease to a much lesser extent when N supply is abundant (Stitt and Krapp, 1999; Fig. 1). Despite the importance of nutrient limitations on carbon fixation, they are frequently ignored in biogeochemical models, including in the case of N (Vitousek and Howarth, 1991; Hungate et al., 2003; Luo et al., 2004; Körner, 2006). For instance, a recent analysis of the Coupled Model Intercomparison Project (CMIP5), one of the most highly regarded multi-model datasets available for predicting future changes in atmospheric CO₂, concluded that CMIP5 has underestimated the atmospheric CO₂ burden projected for the year 2100 by 26–61 ppm, due to a neglect of terrestrial N limitation (Zaehle et al., 2015). Models that, by contrast, incorporate C-N interactions, indicate that the potential for terrestrial C capture could be reduced by 50% or more due to limitations in the nitrogen cycle (Sokolov et al., 2008; Thornton et al., 2009; Arneeth et al.,

2010; Zaehle et al., 2010). In fact, some models even suggest the terrestrial biosphere could turn into a net carbon source by the end of the century (Wieder et al., 2015; Mystakidis et al., 2016), given the numerous factors linked to climate change, including changing precipitation, temperature, ozone levels, microbial interactions, disease risk, and nutrient cycles (Melillo et al., 1993; Cramer et al., 2001; Harvell et al., 2002; Ciais et al., 2005; Hyvönen et al., 2007; Sitch et al., 2007, 2013; Reich et al., 2014).

In this review, we discuss the patterns of eCO₂ on terrestrial plant and ecosystem production, in the context of N limitation, and examine mechanisms occurring at various levels of organization that may explain some of the most pronounced effects. These include N- and eCO₂-dependent changes in growth and yield in grassland, forest, and agricultural systems, N uptake, assimilation, and accumulation under eCO₂, and progressive nitrogen limitation (PNL) and related soil processes.

2. N constraints on plant-growth responses to eCO₂

At the ecosystem level, eCO₂-associated increases in net ecosystem production (NEP, the difference between gross primary production (GPP) and ecosystem respiration, i.e. the sum of heterotrophic and autotrophic respiration) have been observed in many reports. The strongest effects are often seen at higher latitudes (>40°N; Forkel et al., 2016), which is largely attributable to high-latitude warming trends in addition to eCO₂, manifesting as a "greening" trend of increased vegetation cover at such latitudes (Myneni et al., 1997; Lucht et al., 2002). N constraints on soil-ecosystem production are also more prevalent at high latitudes (temperate, boreal, and tundra regions), where biological N₂ fixation (BNF) is naturally low, although the deposition of anthropogenically-derived reactive nitrogen (Nr) can be quite high in these regions (Vitousek and Howarth, 1991; Reich and Oleksyn, 2004; LeBauer and Treseder, 2008; Zaehle, 2013). Coincidentally, the majority of CO₂-N studies at the ecosystem level have been conducted in grassland and forest biomes of higher latitudes, in addition to agricultural systems (Table 1).

In the following discussion, we focus on eCO₂-induced stimulations of NPP in forest, grassland, and agricultural systems (Norby et al., 2005; Luo et al., 2006; Matthews, 2007; Leakey et al., 2009; Franks et al., 2013; cf. Dukes et al., 2005; Inauen et al., 2012), as well as the constraints imposed by N limitations (Reich et al., 2006a; Norby et al., 2010; Reich and Hobbie, 2013; Feng et al., 2015). We also look at soil respiration, which has been shown to consistently increase under eCO₂, and which varies widely in response to varied N levels from one ecosystem to another (Janssens et al., 2010; Adair et al., 2011; Maaroufi et al., 2015; Yue et al., 2016). Because soil respiration releases roughly 10 times more CO₂ into the atmosphere than all combined anthropogenic sources (Schlesinger and Andrews, 2000; Raich et al., 2002), even a modest deviation in soil respiration has the potential to greatly exacerbate or mitigate CO₂ emissions.

2.1. Methodological note

We begin with a brief consideration of methods used to study the effects of eCO₂ on plant carbon fixation. Early investigations relied on growth and physiological analyses conducted using growth chambers, greenhouses, or open-top chambers (OTCs; for review, see Leadley and Drake, 1993; Drake et al., 1997; Curtis and Wang, 1998; Medlyn et al., 1999; Wand et al., 1999; Ainsworth et al., 2002; Jablonski et al., 2002). It has been argued that major limitations arise with such techniques, including size constraints of the chambers, limited growing periods, and difficulties in extrapolating small-scale findings to larger, "real-world", environments




Fig. 1. The CO₂ fertilization effect in relation to N nutrition. Plant biomass increases with available N to a greater extent in plants grown on elevated CO₂ (solid line) relative to ambient CO₂ (dashed line). However, with long-term exposure to elevated CO₂, progressive N limitation (PNL) down-regulates biomass gains, which is more pronounced under low-N conditions (dotted line). Redrawn from Wong, 1979; and modified based on Bowler and Press, 1996; Reich et al., 2006b (cf. Hofmockel et al., 2011; Feng et al., 2015).

(e.g. Ainsworth et al., 2002; Long et al., 2004; Ainsworth and Long, 2005; Bunce, 2012). More recent developments in FACE technology, by contrast, have allowed plants to grow under more natural, fully open-air conditions (Ainsworth and Long, 2005; Leakey et al., 2009). FACE imposes fewer restrictions on plant growth, notably root growth, and is especially well suited to the study of large, long-lived plants, such as trees (Arp, 1991; Curtis and Wang, 1998; McLeod and Long, 1999). However, others have argued that the benefits of FACE may be overstated, and that, as long as strict and realistic consideration of growth conditions are monitored (especially with respect to soil nutrients), techniques using OTCs may be appropriate (Körner, 2006; and references therein). Moreover, FACE technology can be costly (Messerli et al., 2015) and conditions may be difficult to control (such as with N addition). In addition, they often entail significant variations in CO₂ levels in addition to the frequent practice of not providing eCO₂ at night, both of which can have pronounced effects on plant growth (Bunce, 2012, 2014). Nevertheless, because of their scale, duration, and overall fidelity to natural conditions, FACE-type systems have become a preferred means of providing information inputs for models simulating and projecting the effects of eCO₂ on plant growth. Thus, the majority of studies discussed below are ones that have utilized FACE technology (see also Table 1). It may be noted that, as yet, FACE studies are relatively uncommon, in particular those which measure interactions between C and N over decennial timescales (Feng et al., 2015). In addition, FACE studies tend to be geographically biased towards temperate ecosystems of North America and Europe, while much work remains to be done in boreal and tropical regions, which have great significance for C sink and storage activity (Leakey et al., 2012; Jones et al., 2014; Norby et al., 2016), and where climate warming is most pronounced (see above).

The choice among methodologies is not trivial; overall, FACE studies have indicated a substantially lower (in many cases by 50% or more) eCO₂ fertilization effect for a wide variety of crops and herbaceous plants (although, in the case of trees, FACE studies often show a larger effect), with disquieting implications for climate-change mitigation and the future of humanity's food supply (Ainsworth and Long, 2005; Long et al., 2006; Leakey et al., 2009; Weigel and Manderscheid, 2012; cf. Tubiello et al., 2007). It is interesting to note, however, in the context of the present article, that nitrogen status may be more important for their growth responses

to eCO₂ than the experimental system used (Yin, 2013; Pleijel and Högy, 2015).

2.2. Grasslands

Multiple studies have indicated that C₃ grasses respond positively to eCO₂ in terms of biomass gain, and that this response is frequently N-limited. For instance, in a study of 13 grassland species over 11 years at the BioCON (Biodiversity, CO₂, and Nitrogen) FACE site (Table 1), it was shown that net photosynthesis (A_{net}) was stimulated by about 10% with eCO₂ (Lee et al., 2011). Superimposed on this was a strong and persistent N limitation on eCO₂-induced biomass gain, as indicated by a marked reduction in growth stimulation in plots lacking N fertilization (Reich et al., 2006a; Reich and Hobbie, 2013; Figs. 1 and 3). Intriguingly, however, no significant CO₂ × N interactions were observed in the majority of the response variables tested: A_{net} , specific leaf area (SLA), stomatal conductance (g_s), instantaneous water-use efficiency, or photosynthetic N-use efficiency (PNUE) (Lee et al., 2011). In another seminal long-term grassland study, the SwissFACE experiment investigated CO₂ × N effects in a perennial ryegrass (*Lolium perenne*) sward, with similar results. In this setting, Schneider et al. (2004) observed an increased harvestable biomass (+32%) under eCO₂ but only at high N supply (56 g N m⁻² yr⁻¹ vs. 14 g N m⁻² yr⁻¹). An earlier study in the SwissFACE plot showed no significant CO₂ × N interaction in daily integrated CO₂ uptake (A'), light-saturated CO₂ uptake (A_{sat}), or g_s (Ainsworth et al., 2003); however, low-N plants exhibited twice the extent of photosynthetic acclimation (down-regulation) over the course of long-term eCO₂ application (see Section 3.1; Figs. 2 and 3). The question of whether leaf-level CO₂ × N interactions (especially those pertaining to photosynthetic activity and capacity) scale to overall plant production (NPP), and further to ecosystem carbon capture (NEP), remains a subject of considerable debate (Reich et al., 2006b; Fernández-Martínez et al., 2014; Körner, 2015). As pointed out by Reich et al. (2006b), C-N interactions may manifest less prominently at the small (leaf) scale than at the large (ecosystem-wide) scale, due to the added effects of growth and decay dynamics of root systems and tree canopies. Körner (2015) has argued that the widespread assumption that photosynthesis is the rate-controlling factor of plant growth, may be mistaken;




Fig. 2. Photosynthetic response to CO_2 enrichment and available N. (A) CO_2 assimilation rate, measured at growth [CO_2] (ambient or elevated; dashed and solid lines, respectively), as a function of available N. (B) CO_2 assimilation rate, measured at ambient [CO_2], as a function of available N. Note, photosynthetic acclimation occurs in plants grown on elevated CO_2 and is most pronounced under low-N conditions. Redrawn from Wong, 1979 (see also Sage, 1994; Bowler and Press, 1996; Miglietta et al., 1996; Stitt and Krapp, 1999; cf. Farage et al., 1998).

instead, it was proposed that sink capacity (tissue growth) controls C fixation based on demand.

Biomass gains with eCO_2 are not ubiquitous in grasslands, however. Shaw et al. (2002) showed that plants indeed responded to eCO_2 in terms of increased NPP in a California grassland, but only when eCO_2 was used as a single-factor treatment. Multi-factor combinations of eCO_2 and increased temperature, precipitation, and N deposition, on the other hand, actually decreased NPP, relative to ambient CO_2 conditions. The authors hypothesized that other soil nutrient limitations may be at play, perhaps due to gradual immobilization of nutrients by soil microorganisms, or decreased root allocation, which would limit the capacity of plants to forage for nutrients. Similarly, Dukes et al. (2005) found no biomass gains with eCO_2 , irrespective of N supply, and suggested that phosphorus (P) limitations may be central to this observation, because eCO_2 reduced total plant P uptake. Inauen et al. (2012)

also showed no growth stimulation by eCO_2 irrespective of N fertilization in nine species of plants collected from a glacier forefield (secondarily successional bare land, exposed by glacial retreat). However, they did observe a drastic change in root-shoot biomass partitioning, with aboveground biomass reduced by 35% across all species, and higher biomass allocated to the roots. The authors suggested that these alpine plants were not carbon-limited under current conditions. It is worth noting as well that land-use intensity (e.g. livestock grazing, fertilizer usage, etc.) can be a dominant driver of soil C sequestration in grasslands (Chang et al., 2016). Overall, these findings emphasize the need for a better understanding of soil nutrient constraints on plant and ecosystem production under eCO_2 .

Ecosystem respiration is consistently stimulated in response to eCO_2 in grasslands (Luo et al., 1996; Zak et al., 2000; Craine et al., 2001; Adair et al., 2011). In a dramatic example, Adair et al. (2011)

Table 1
Survey of studies examining CO₂ × N interactions on biomass and photosynthetic response.

Study/System	Species/Ecosystem	Age/Time frame	Treatments	Biomass response	Photosynthetic response	References
Swiss FACE	<i>Lolium perenne</i> L. swards	10 years	360 and 600 ppm CO ₂ , 14 and 56 g NH ₄ NO ₃ m ⁻² yr ⁻¹	Increased CO ₂ effect with high N (decline over time).	No effect on A _{sat} , A', g _s , or J _{max} . Significant decrease in V _{c,max} in low-N plants.	Ainsworth et al. (2003) and Schneider et al. (2004)
BioCON FACE	Perennial grassland (legumes, non-leguminous forbs, C ₃ and C ₄ grasses)	5–10 years	360 and 560 ppm CO ₂ , ambient and +4 g NH ₄ NO ₃ m ⁻² yr ⁻¹	Increased CO ₂ effect with high N after 4–6 years.	No effect on A _{net} , A _{sat} , SLA, or g _s .	Lee et al. (2001), Reich et al. (2006a,b), Lee et al. (2011), Reich and Hobbie (2013) and Reich et al. (2014)
Oak Ridge National Laboratory (ORNL) FACE	Sweetgum (<i>Liquidambar styraciflua</i>)	11 years	Ambient and 550 ppm CO ₂ , ambient and +200 kg urea-N ha ⁻¹ yr ⁻¹	Stimulation in wood increment with N fertilization and eCO ₂ .	N/A	Norby et al. (2005, 2010)
Duke FACE	Loblolly pine (<i>Pinus taeda</i>)	10 years	Ambient and 550 ppm CO ₂ , ambient and +11 g urea-N m ⁻² yr ⁻¹	Increased NPP with eCO ₂ and N fertilization.	N/A	Finzi et al. (2007), McCarthy et al. (2010) and Drake et al. (2011)
Jasper Ridge Global Change Experiment	Annual grasses (<i>A. barbata</i> , <i>B. hordeaceus</i>) and forbs (<i>G. dissectum</i> , <i>E. botrys</i>)	3–5 years	Ambient and 680 ppm CO ₂ , ambient and +3.5 g Ca(NO ₃) ₂ m ⁻² yr ⁻¹	Reduced NPP under eCO ₂ compared to ambient CO ₂ .	N/A	Shaw et al. (2002), Zavaleta et al. (2003) and Dukes et al. (2005)
Cedar Creek FACE	Herbaceous brackish wetland (C ₃ sedge, C ₄ grasses)	4 years	Ambient and 720 ppm CO ₂ , ambient and +25 g NH ₄ Cl m ⁻² yr ⁻¹	eCO ₂ -induced increase in C ₃ biomass and decrease in C ₄ biomass with added N.	N/A	Langley and Magonigal (2010)
Japan Rice FACE	<i>Oryza sativa</i> L.	3 years	365 and 600 ppm CO ₂ , low (4), medium (8–9), and high (12–15 g m ⁻²) N (as (NH ₄) ₂ SO ₄)	c. 15% increase in grain yield with eCO ₂ for medium- and high-N crops; no effect on total biomass; strong dependence on developmental stage.	N/A	Okada et al. (2001), Kim et al. (2003) and Shimono et al. (2008)
MeteoSwiss FACE/Mini FACE	Glacier forefield plants (graminoids, forbs, cushion)	3 years	385 and 580 ppm CO ₂ , ambient and +25 kg N ha ⁻¹ yr ⁻¹	Aboveground biomass decline with eCO ₂ , irrespective of N supply.	Decline in A _{net} .	Inauen et al. (2012)
Wheat MiniFACE	Winter wheat	1 year	Ambient and 600 ppm CO ₂ , ambient and +15.6 g urea m ⁻²	N/A	Significant decrease in A _{net} , g _s , V _{c,max} and J _{max} under eCO ₂ and low N.	Miglietta et al. (1996)
Swiss model ecosystem OTC	Beech (<i>Fagus sylvatica</i> L.) and Norway spruce (<i>Picea abies</i> Karst.)	5 years	370 and 570 ppm CO ₂ , 0.5 and 5 g NH ₄ NO ₃ m ⁻² yr ⁻¹	Increased NPP.	N/A	Spinnler et al. (2002)
Chamber, pots	<i>Bromus mollis</i>	130 d	350 and 650 ppm CO ₂ , 0.37 and 1.87 mM NH ₄ NO ₃	Increased seed biomass.	No effect on A _{net} .	Larigauderie et al. (1988)
Chamber, hydroponics	<i>Beta vulgaris</i> L.	7-weeks old	32 and 100 Pa CO ₂ , 7.5 mM NO ₃ ⁻ or NH ₄ ⁺	Increase biomass with NO ₃ ⁻ .	Increased A _{net} in NH ₄ ⁺ -grown plants relative to NO ₃ ⁻ -grown plants.	Raab and Terry (1994)

eCO₂ = elevated CO₂; A' = daily integrated CO₂ uptake; A_{sat} = light-saturated CO₂ uptake rate; A_{net} = net photosynthesis; g_s = stomatal conductance; V_{c,max} = maximum velocity of Rubisco carboxylation; J_{max} = maximum rate of electron transport; SLA = specific leaf area.

found that eCO₂ increased soil respiration by 45% in the BioCON plot. This large effect was found to be the result of increased C inputs and availability, rather than a result of changing soil water availability (eCO₂ has been found to reduce stomatal conductance and plant water use, and thereby increase soil moisture; Adair et al., 2011). However, unlike plant biomass production, soil respiration was found not to be strongly N-limited in BioCON (Adair et al., 2011). Unfortunately, there is currently a lack of similar multi-factor studies proceeding over a longer time frame, particularly with respect to soil respiration in grasslands. It is critical that such studies be

forthcoming, given the major impact soil C uptake and release has on the global C budget (Raich et al., 2002; Le Quéré et al., 2015).

2.3. Forests

Among terrestrial plants, trees are exceptionally strong “beneficiaries” of eCO₂, particularly when young. FACE studies have demonstrated increases in biomass approaching or even exceeding 50% under eCO₂ (Dijkstra et al., 2002; Ainsworth and Long, 2005; Ainsworth and Rogers, 2007). Thus, forests are an especially

important terrestrial C sink due to their capacity to sequester large amounts of C from the atmosphere into long-lived biomass and into soil organic matter (SOM) *via* litter. Globally, forests cover more than 4 billion hectares of land area and possess an estimated total C-sink capacity of $2.4 \pm 0.4 \text{ Pg C yr}^{-1}$ (Pan et al., 2011). Although eCO_2 increases forest NPP (Norby et al., 2005; Hyvönen et al., 2007), soil N limitations can impose a major growth constraint, as in other ecosystems (Norby et al., 2010). For example, in an 11-year FACE experiment in a sweetgum (*Liquidambar styraciflua*) forest stand, eCO_2 -induced NPP stimulation declined from 24% to 9% over the final 5 years of the study. This was attributed to the decline in soil N availability under eCO_2 , as more N was sequestered into wood and SOM (an example of “progressive nitrogen limitation” (PNL); see Section 3.4), thus limiting tree growth (Norby et al., 2010). By contrast, PNL does not seem to have occurred in a decade-long FACE experiment with loblolly pine (*Pinus taeda*) in the Duke FACE plot (Finzi et al., 2007; Hofmockel et al., 2011). It is suggested that increased N acquisition (through some combination of increased fine root production, SOM decomposition, and C allocation to mycorrhizal associations) sustains the increases in NPP under eCO_2 (Finzi et al., 2007; Drake et al., 2011; Hofmockel et al., 2011; Phillips et al., 2011; see also Sections 3.3 and 3.4).

In general, forest production benefits from anthropogenic N deposition, and in several cases, soil N has been found to be the primary determinant of NEP in forested systems (Magnani et al., 2007; Norby et al., 2010; Fernández-Martínez et al., 2014; Maaroufi et al., 2015; Yue et al., 2016). In a study of 92 forests stands, including boreal, temperate, Mediterranean, and tropical forests, Fernández-Martínez et al. (2014) determined that only nutrient-rich forests increase carbon sequestration (NEP) *via* increasing carbon uptake. Ecosystem carbon-use efficiency (CUE_e, i.e. the ratio of NEP to GPP) in nutrient-rich forests was found to be more than 5-fold higher than that in nutrient-deficient forests (33% *versus* 6%, respectively). The conclusions of this study, however, while striking, are in conflict with the great majority of global coupled carbon-climate models, which assume that C inputs *via* photosynthesis drive plant production and C sequestration. Rather, it emphasized that nutrient limitation is the single most important determinant of forest carbon sequestration globally (see Section 2.2; Körner, 2015).

The much lower CUE_e in nutrient-poor forests observed in the Fernández-Martínez et al. (2014) study was attributed to a much larger proportion of GPP being lost to ecosystem respiration. The authors proposed that limited nutrient availability decreases C allocation to long-lived woody tissues, instead increasing C allocation to fungal root symbionts and root exudates, thus stimulating heterotrophic respiration in the rhizosphere (Janssens et al., 2010; Vicca et al., 2012), and stimulating overall ecosystem respiration. Conversely, several studies have shown that increased N deposition can mitigate the eCO_2 -induced stimulation of soil respiration (Burton et al., 2004; Olsson et al., 2005; Janssens et al., 2010; Maaroufi et al., 2015; see also Fig. 3). However, exceptions do exist. For instance, Hasselquist et al. (2012) found that moderate N addition rates, similar to typical N deposition rates in boreal regions ($20 \text{ kg N ha}^{-1} \text{ yr}^{-1}$), resulted in a doubling of soil respiration compared to control plots. This enhanced respiration was linked to increased autotrophic respiration. This study highlights the need for more investigation into $\text{CO}_2 \times \text{N}$ interactions in soil respiration, particularly exploring a range of realistic N deposition rates (as opposed to simple binary treatments).

Despite having the highest rates of NPP on Earth ($\sim 10 \text{ Pg C yr}^{-1}$; 30–40% of global NPP; Cleveland et al., 2015), tropical forests have been inadequately studied in terms of eCO_2 effects and nutrient limitation (Cleveland et al., 2011, 2015; Norby et al., 2016). In a recent meta-analysis of 113 tropical forest sites, Cleveland et al. (2011) suggest that P availability may be the most limiting nutrient to NPP in such systems; however many more studies, including

those employing FACE (Norby et al., 2016), will be necessary to establish the chief limiting nutrients in different forest ecosystems (see also Vitousek, 1984; Wieder et al., 2015). A joint focus on N and P is clearly necessary.

2.4. Agriculture

As with natural ecosystems, many studies have shown that the productivity of agricultural C_3 plants, including major crops such as rice, wheat, potato, soybean, cotton, and tobacco, can be significantly increased by eCO_2 , and such increases also depend on N availability in most cases (Kimball et al., 2002; Kim et al., 2003; Ainsworth and Long, 2005; Sakai et al., 2006; Ziska and Bunce, 2007; Fig. 3). For example, in a FACE experiment with wheat, Ma et al. (2007) found that the eCO_2 -stimulated biomass under high N (250 kg N ha^{-1}) was approximately double that under low N (125 kg N ha^{-1}). In a survey of studies using FACE and earlier chamber-based methods, Kimball et al. (2002) showed that eCO_2 stimulated the growth of C_3 plants (wheat, perennial ryegrass, rice, white clover, potato, cotton, and grape) by 17%, on average, when N was adequately supplied, but only by 3% under low N (when the legume was excluded; see below); only in rare cases were exceptions to this general result found, such as in increases in the yield of perennial ryegrass, under low compared to high N.

By contrast, in a six-year FACE study of a crop rotation with winter barley, ryegrass, sugar beet, and winter wheat, Weigel and Manderscheid (2012) found that, contrary to expectations, eCO_2 effects on plant growth variables were independent of N supply. The authors noted that, in a handful of prior studies in rice, grain yield was also independent of N supply (Liu et al., 2008; Yang et al., 2009), although in one of these studies this independence only occurred between medium- and high-N provision, while an eCO_2 -driven yield increase was still found between low- and medium-N treatments (Kim et al., 2003). In rice, yield stimulations or reductions in response to eCO_2 appear to be cultivar-dependent (Satapathy et al., 2014).

More broadly, the magnitude of crop responses to eCO_2 varies widely with the type of plant examined. Among C_3 plants, those with large C sinks in the form of carbohydrate storage organs, such as potato and cassava, and woody tissues such as cotton, can have particularly strong growth responses to eCO_2 (Ainsworth and Long, 2005; Rosenthal et al., 2012). In cassava, a 100% yield increase has been observed under eCO_2 with N fertilization (Rosenthal et al., 2012), and even with low N inputs tuber mass was shown to increase by 49% (Fernández et al., 2002). Because photosynthetic and growth responses of C_4 plants to eCO_2 are generally much lower than those of C_3 plants (Ainsworth and Long, 2005; cf. Wand et al., 1999), agricultural productivity increases for major C_4 crops such as corn, sugarcane, sorghum, and millet, may be limited in a CO_2 -enriched atmosphere, except under drought conditions (Lopes et al., 2011). By contrast, positive eCO_2 effects on growth and yield in legume species tend to be particularly pronounced, due to their ability to fix organic N from atmospheric N_2 *via* root symbioses (Zanetti et al., 1996; Ainsworth et al., 2002; Rogers et al., 2006; Kant et al., 2012). However, whereas they may overcome N limitation in this way, legumes are often phosphorus limited (Rogers et al., 2009; see below), underscoring our earlier point on the necessity to examine not only N-, but also P-limitation effects under eCO_2 .

A potential threat to agricultural production as C_a continues to rise is the reduction of N concentrations in tissues of crop plants, associated with increased biomass gains (see Section 3.1). In cereals, this is reflected in an 8–31% reduction in grain N, mostly in the form of organic N, i.e. protein and amino acids (Fangmeier et al., 1999; Kant et al., 2012; Pleijel and Uddling, 2012). The reduced protein levels in cereal grains can be gravely detrimental for human

nutrition (Taub et al., 2008), as well as for the quality of bread flour produced from these grains (Högy et al., 2009).

3. Plant and soil N uptake and accumulation under eCO₂

3.1. Tissue N content

Many studies have indicated that plants grown under eCO₂ show decreased tissue concentrations of N relative to plants grown under ambient CO₂, although this condition can be alleviated by increased N supply, uptake, and -use efficiency (Stitt and Krapp, 1999; Kant et al., 2012; Fig. 3). The decrease in the concentration of N in plant dry mass (N_m) varies between studies, but typically falls in the range of 9–16% depending on plant organ, species, functional group, or experimental method (Cotrufo et al., 1998; Curtis and Wang, 1998; Ainsworth and Long, 2005; Taub and Wang, 2008; Feng et al., 2015). Understanding the nature of this decrease is critical to the development of comprehensive models of carbon fixation, plant growth and yield, and ecosystem-level carbon sequestration (NEP) under eCO₂. Although the underlying mechanisms responsible for the decrease in N_m under eCO₂ are not fully understood, several hypotheses have been put forward (Taub and Wang, 2008; Bloom, 2015a; Feng et al., 2015).

The hypothesis most often put forward is that of ‘biomass dilution’ of tissue N_m , in response to increased photosynthetic assimilation of C under eCO₂ (Luo et al., 1994; Poorter et al., 1997; Farage et al., 1998; Gifford et al., 2000; Yin, 2002; Ellsworth et al., 2004; Taub and Wang, 2008; cf. Feng et al., 2015). Biomass dilution occurs when the biomass gains (NPP) under eCO₂ are greater than the corresponding changes in N uptake (see Section 3.3; it should be noted here that tissue N content per unit leaf area also declines; Ellsworth et al., 2004). However, this appears to be only partly responsible for the decrease in N_m . In a recent meta-analysis of studies in grassland, forest, and cropland ecosystems, Feng et al. (2015) asserted that the change in N_m in response to eCO₂ was in fact independent of the biomass (NPP) response to eCO₂. The authors suggest that the N limitations imposed on plant NPP gains under eCO₂ (and the corresponding decreases in N_m) are more likely to be linked to eCO₂-induced suppressions of plant N acquisition (Section 3.3), rather than to growth dilution of plant N or to processes related to progressive N limitation (Section 3.4). A further argument against the biomass dilution hypothesis is the disproportionate decrease in tissue N compared to decreases in tissue levels of other nutrients (e.g. potassium (K) and P; Poorter et al., 1997; Cortufo et al., 1998; Taub and Wang, 2008; Jauregui et al., 2016), suggesting that processes other than biomass dilution are affecting N_m .

The well characterized decreases in ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) content under eCO₂ may account for some of the decreases observed in N_m (Long et al., 2004; McMurtrie et al., 2008; Taub and Wang, 2008). Approximately 25–30% of total leaf N is found in Rubisco, which is widely accepted to be the rate-limiting step of photosynthesis (Sage et al., 1987; Jensen, 2000; Spreitzer and Salvucci, 2002). With the increase in carboxylation efficiency under eCO₂ (due to the higher ratio of CO₂ to O₂ at the active site of Rubisco, favoring photosynthesis over photorespiration), C supply can exceed demand, particularly when N limitations constrain sink development; thus, it has been hypothesized that N from Rubisco can be recycled to offset limitations in sink development (Drake et al., 1997; Rogers et al., 1998; Long et al., 2004; McMurtrie et al., 2008). According to Moore et al. (1999), the reduced expression of genes encoding Rubisco are brought about by the hexokinase-mediated signaling of increased photosynthate levels. The eCO₂-induced declines in Rubisco are considered the basis of the acclimation response in which photosynthetic capacity is down-regulated under eCO₂, particularly under N-limiting conditions (Wong, 1979; Oberbauer

et al., 1986; Arp, 1991; Sage, 1994; Drake et al., 1997; Rogers et al., 1998; Stitt and Krapp, 1999; Ainsworth and Rogers, 2007; see also Figs. 2 B and 3). However, it is worth noting that even when Rubisco levels are down-regulated under acclimation, eCO₂ still increases NPP in most cases (Leakey et al., 2009).

3.2. The effects of contrasting N sources

The relative proportions, and concentrations, of the major N sources available to plants, i.e. nitrate (NO₃⁻), ammonium (NH₄⁺), and organic N (e.g. amino acids), can vary substantially, and this can have profound effects on species distribution and plant performance in the field (Gigon and Rorison, 1972; Haynes and Goh, 1978; Kaye and Hart, 1997; Kronzucker et al., 1997; Hofmockel et al., 2007, 2010; Bloom, 2015b; Rubio-Asensio et al., 2015). Generally, low pH, low temperature, the accumulation of phenolic-based allelopathic compounds, and low oxygen supply favour net ammonification over nitrification in soils (Vitousek et al., 1982; Eviner and Chapin, 1997; Britto and Kronzucker, 2002). For example, late-successional forest soils are typically NH₄⁺-dominated, in contrast to early-successional or post-disturbance forest soils, which are generally NO₃⁻-dominated (Smith et al., 1968; Rice and Pancholy, 1972; Klingensmith and Van Cleve, 1993; Kronzucker et al., 1997). Moreover, periodically flooded agricultural systems like rice paddies and cranberry fields are predominately NH₄⁺-rich (Britto and Kronzucker, 2002; and references therein; see, however, Kirk and Kronzucker, 2005).

Plant responses to eCO₂ can be influenced by N source, although studies examining this are sparse and varied. In sugar beet (*Beta vulgaris*), net photosynthesis was substantially higher at elevated CO₂ (100 Pa vs. an ambient level of 32 Pa) in NH₄⁺-grown plants compared to NO₃⁻-grown plants, due to an increase in photosynthetic enzymes, including Rubisco (Raab and Terry, 1994). Moreover, NH₄⁺ plants showed a doubling of chloroplast volume, a ~60% rise in chlorophyll content, and a 4.3-fold higher accumulation of soluble proteins (see also Guo et al., 2007; for review). In cassava (*Manihot esculenta*), NO₃⁻-grown plants experienced significant photosynthetic acclimation, as opposed to NH₄⁺-grown plants, which showed none; however, this was only the case in young (36-d-old) plants, but was not observed at more advanced ages, at which plants experienced NH₄⁺-toxicity symptoms including severely reduced biomass (Cruz et al., 2014). By contrast, in two grass species, colonial bent (*Agrostis capillaris*) and matgrass (*Nardus stricta*), no significant effect of N form (NH₄⁺ vs. NO₃⁻) was found in any of the measured parameters under eCO₂: total biomass, carboxylation efficiency, A_{max} , leaf N content, specific leaf area, leaf area ratio, or soluble carbohydrate content (Bowler and Press, 1996).

It has been reported that NH₄⁺ supply increases the rate of non-photosynthetic (“dark”) carbon fixation in roots, in contrast to NO₃⁻ supply (Ikeda et al., 1992; Cramer et al., 1993; Koga and Ikeda, 2000; Balkos et al., 2010). This may be because, at elevated NO₃⁻ supply, NO₃⁻ that is not assimilated can be safely stored in root vacuoles (Salsac et al., 1987; Sechley et al., 1992), while, by contrast, NH₄⁺ must be reduced immediately to avoid cytotoxicity (Givan, 1979; Oaks, 1994; Britto and Kronzucker, 2002). Products of dark carbon fixation, mediated mainly by phosphoenolpyruvate carboxylase (PEPC), can anapleuratically provide the carbon skeletons necessary for NH₄⁺ assimilation into organic N (Chollet et al., 1996; Huppe and Turpin, 1994). In fact, when NH₄⁺ is supplied, PEPC activity and expression in roots has been shown to be significantly enhanced (Schweizer and Erismann, 1985; Arnozis et al., 1988; Koga and Ikeda, 2000; Balkos et al., 2010; Britto et al., 2014). In wheat seedlings, treatment with methionine sulfoximine (MSX), an inhibitor of glutamine synthetase (the enzyme that catalyzes primary NH₄⁺ assimilation), suppressed CO₂ fixation in roots, sug-

gesting a role for the products of NH_4^+ assimilation in the regulation of dark fixation (Koga and Ikeda, 2000). Although feeding plants with NH_4^+ may seem like a promising means of increasing CO_2 fixation under eCO_2 , in tomato roots PEPC activity was shown to be suppressed by increased root-zone CO_2 , despite being stimulated by NH_4^+ supply (Viktor and Cramer, 2005).

Bloom et al. (2010) reported that nitrate reductase (NR) activity in both *Arabidopsis* and wheat is limited under eCO_2 . This effect was attributed to the decrease of photorespiration (which can stimulate the supply of reductant for NO_3^- reduction) under this condition, and also to the inhibition of nitrite (NO_2^-) transport from the cytosol to the chloroplast by elevated levels of bicarbonate ions in the leaf. The authors suggested that these effects may explain the differential responses observed in wheat and *Arabidopsis* to eCO_2 under contrasting N sources, where growth enhancement doubled under ammonium supply relative to NO_3^- supply, and shoot protein content declined in NH_4^+ -grown plants to only half the extent observed in NO_3^- -grown plants (Bloom et al., 2002, 2010). Moreover, it may explain the decline in plant NO_3^- acquisition and increase in soil NO_3^- availability, under eCO_2 , and the lack of corresponding responses under NH_4^+ supply, as shown in a recent meta-analysis covering grasslands, forests, and croplands (Cheng et al., 2012). The decrease in organic N circulating from shoot to root as a result of decreased NR activity, however, may signal the root to, in fact, increase NO_3^- uptake (Imsande and Touraine, 1994; Jackson et al., 2008), as was observed by Jauregui et al. (2016), who found that N accumulation, and expression of several NO_3^- transporters in the root, were enhanced by eCO_2 , while shoot NR activity was reduced. However, while the conclusions of Bloom et al. (2010) are based on converging results from five independent methods, some previous studies have indicated that NO_3^- assimilation does not decrease under eCO_2 except when NO_3^- supply is limiting (Nunes-Nesi et al., 2010). For example, several studies have shown a large stimulation of NR activity in tobacco plants under eCO_2 when NO_3^- provision was high, but not when it was low or when NH_4^+ was the N source (Geiger et al., 1998, 1999). Moreover, Kaiser and Forster (1989) showed that NO_3^- reduction was decreased under low CO_2 concentrations, and that this inhibition was reversed when CO_2 was added back to the system. More investigations are clearly warranted to address the apparent discrepancies in this complex interaction.

3.3. Plant N acquisition

The effects of eCO_2 on plant N acquisition are very likely to contribute to changes in plant N_m ; however, N-acquisition responses to eCO_2 can be quite variable. Feng et al. (2015) observed a positive, linear, and strong relationship between the effect of eCO_2 on N acquisition and the effect of eCO_2 on plant production (in contrast to the response of N_m ; see above). Interestingly, however, they also showed that N acquisition under eCO_2 decreased by up to ~10% relative to ambient- CO_2 levels when the corresponding plant production gains were nil or only modestly positive, regardless of N supply. According to the authors, this was likely due to plants having decreased access to N under these conditions.

Several hypotheses have been put forward to explain the decrease in N acquisition, where it occurs (e.g. Taub and Wang, 2008; Feng et al., 2015), including: (i) slower decomposition of plant litter with high C:N ratios, thus decreasing soil N availability over time (Luo et al., 2004), (ii) net N immobilization by a larger soil microbial biomass (de Graaff et al., 2006), (iii) shoot-to-root signaling of reduced aboveground N demand, either via inhibition of nitrate (NO_3^-) assimilation (Bloom et al., 2010; Bloom, 2015a; Jauregui et al., 2016) or decreased abundance of photosynthetic enzymes (especially Rubisco; Long et al., 2004; McMurtrie et al., 2008), (iv) decreased transpiration-driven mass flow of soil N to roots (McDonald et al., 2002; Cramer et al., 2009; McGrath

and Lobell, 2013; Matimati et al., 2014), (v) decreased expression levels of root N transporters, such as NRT1.1. (Guo et al., 2013), and (vi) altered root system architecture, including increases in fine and coarse root mass, and root:shoot ratios (Beidler et al., 2015). By contrast, hypotheses for the stimulation in N acquisition include: (i) enhanced foraging by roots and increased whole-plant N sink strength (Finzi et al., 2007; Iversen et al., 2012), (ii) stimulated decomposition and N mineralization by a larger and/or more active microbial community (Dijkstra et al., 2008; Iversen et al., 2011; Phillips et al., 2011), (iii) increased biological N_2 fixation (Poorter et al., 1997; Rütting and Andresen, 2015; cf. Hungate et al., 2004), and (iv) increased N transporter activity and/or expression (Jauregui et al., 2016).

Despite the general observation of reduced N concentrations in plant tissues under eCO_2 , however, N uptake by roots nevertheless appears to be enhanced under these conditions in many cases, although the extent of this enhancement is less than the increase in C fixation (Stitt and Krapp, 1999; Johnson, 2006; Lenka and Lal, 2012). Several earlier reviews reported no consistent pattern in terms of changes in the direction or extent of root nutrient absorption capacity under eCO_2 , possibly due to variable experimental protocols and differences among species (e.g. BassiriRad et al., 2001), or the extent of N supply (Stitt and Krapp, 1999). More recent studies and reviews have generally indicated positive effects of eCO_2 on N uptake (Torbert et al., 2004; Yamakawa et al., 2004; Johnson, 2006; Finzi et al., 2007; Zak et al., 2007; Rennenberg et al., 2010; Hofmockel et al., 2011), with at times very large increases, for example a 33–34% greater total N in the case of sorghum and soybean (Torbert et al., 2004). Mechanistically, studies have tended to focus on changes in N uptake as a function of changes in soil-N availability, root-system size and architecture, and mass flow (e.g. Finzi et al., 2007; Hofmockel et al., 2011; McGrath and Lobell, 2013).

Relatively little work has examined changes in N acquisition under eCO_2 at the cellular or molecular level. Matt et al. (2001a,b) showed that, in tobacco, altered diurnal expression patterns of the genes encoding nitrate reductase and the nitrate transporter NRT2 resulted in overall higher N uptake and assimilation with NO_3^- as the N source (cf. Bloom et al., 2010). When grown on a mixed-N source (NH_4NO_3), however, the plants preferentially took up NH_4^+ , which was suggested to be due to a large capacity for NH_4^+ assimilation brought about by eCO_2 and its release of photorespiratory demands on NH_4^+ re-assimilation (Raven et al., 1992; see also Section 3.2). In neither case was a reduction of tissue N concentration seen, indicating the presence of an adequate N supply (2 mM). In another example, yellow birch (*Betula alleghaniensis*) showed NO_3^- uptake was reduced and NH_4^+ uptake was unchanged with eCO_2 , despite significant increases in total root length being observed in both conditions (Bauer and Bernston, 2001). By contrast, in the same study, no change in root system architecture with eCO_2 was observed in white pine (*Pinus strobus*), whereas both NH_4^+ and NO_3^- uptake were stimulated. A recent study by Jauregui et al. (2016) demonstrated large increases in the expression of genes encoding NO_3^- transporters in *Arabidopsis*, coinciding with a doubling of NO_3^- concentration in roots. By contrast, Guo et al. (2013) found that eCO_2 downregulated nitrate transport and transporter expression in the legume *Medicago truncatula*. Importantly, however, a proliferation of root nodules was also observed, with concomitant increases in expression of genes encoding N-fixation-related proteins as well as N fixation itself, resulting in higher tissue-N concentrations under eCO_2 .

3.4. Progressive N limitation (PNL) and N deposition

The effects of eCO_2 on soil N levels and cycling have been highly variable, having been shown to decrease, remain stable, or increase under eCO_2 (Gill et al., 2002; Finzi and Schlesinger, 2003; Luo et al.,

2004; Schneider et al., 2004; Reich et al., 2006a,b; Dijkstra et al., 2008; Langley et al., 2009; Phillips et al., 2009; Norby et al., 2010; Zak et al., 2011; Mueller et al., 2013; Rütting and Andresen, 2015).

The progressive nitrogen limitation (PNL) hypothesis proposes that long-term exposure to eCO_2 may result in feedbacks that suppress the availability of N, which in turn reduces the CO_2 fertilization effect (Luo et al., 2004; Dukes et al., 2005; Hungate et al., 2006; Reich et al., 2006b). This effect is particularly pronounced when N supply is low at the outset (Stitt and Krapp, 1999; Figs. 1 and 3). More specifically, PNL postulates that soil N availability becomes increasingly limited as it is sequestered into long-lived plant biomass (such as woody tissues) and recalcitrant fractions of SOM (Luo et al., 2004). In what is perhaps the most striking example of PNL, the decline in soil N availability was linked to the decline in eCO_2 -stimulated NPP in the Oak Ridge National Laboratory (ORNL) FACE experiments (Norby et al., 2010; Norby and Zak, 2011; see Section 2.3). A similar conclusion was drawn in two grassland ecosystems (Reich et al., 2006a; Hovenden et al., 2008). By contrast, little evidence for PNL, even in the absence of N fertilization, has been found in the Duke FACE experiment, where loblolly pine has been growing at eCO_2 -stimulated rates for over a decade (Finzi et al., 2007; McCarthy et al., 2010; Hofmockel et al., 2011; see below). In this system, it appears that PNL may be overcome by a combination of increased growth of deep roots, increased mineralization of N, and a shift in mycorrhizal fungal distribution towards deeper soil (Pritchard et al., 2008; Hofmockel et al., 2011). It is important to note, as Lenka and Lal (2012) point out, that turnover times of many of the recalcitrant biomass and SOM pools in question are typically much longer (e.g. >25 years; Parton et al., 1987) than the length of most FACE experiments (5–10 years), and, thus, much longer experiments are needed to fully understand the effects on soil nutrient cycles. Feng et al. (2015) even suggest that PNL may not be commonly occurring at all, since the effects of eCO_2 on plant production and N acquisition do not diminish over the decennial timescale in their meta-analysis.

Dieleman et al. (2012) caution that many PNL-related studies are based on single-factor treatments (i.e. eCO_2) and results may vary depending on co-factors. One critical factor in this context is soil warming. For example, when warming temperatures were considered in the meta-analysis of Dieleman et al. (2012), soil-N levels were found not to decline to the same extent as with eCO_2 alone,

possibly due to warming-induced accelerations in litter decomposition (see also Melillo et al., 2011; Davidson and Janssens, 2006; Hovenden et al., 2008; Bai et al., 2013). In another study, soil warming ($5^\circ C$ above ambient) for 7 years in a deciduous forest resulted in an average 45% annual increase in net nitrogen mineralization and a three-fold increase in nitrification since the start of the experiment (Butler et al., 2012). The warming-induced increases in bioavailable soil N resulted in significant increases in growth rate and foliar N content in the surrounding trees, especially red maples. Although it was not observed in the study, the increased availability and proportion of soil NO_3^- suggest that species adapted to this N source may be at a significant competitive advantage over NH_4^+ -preferring species. Overall, these studies further highlight the importance of considering the multiple, and at times, contrasting, factors affecting ecosystem processes under eCO_2 and associated climate change.

Anthropogenic N deposition has had significant effects on the C cycle and plant carbon fixation, and is often recommended under PNL conditions; however, its net contribution to climate change remains widely debated (Galloway et al., 2008; Meyerholt and Zaehle, 2015). The extent of N deposition has increased tremendously since pre-industrial times, due to nitrogenous emissions from fossil fuel combustion and the industrial production of Nr species via the Haber-Bosch process, which consumes up to 5% of the world's natural gas (about 1% of total global energy supply), thus itself contributing to eCO_2 (Schlesinger, 2000; Smith, 2002; Smil, 2004). N deposition has increased from $\sim 15 \text{ Tg N yr}^{-1}$ in 1860 to $\sim 187 \text{ Tg N yr}^{-1}$ by 2005 (more than a 12-fold increase), and is projected to rise to $\sim 267 \text{ Tg N yr}^{-1}$ by 2050 (Galloway et al., 2004, 2008). Unfortunately, N-use efficiency by plants has also declined pervasively, resulting in a cascade of environmental and human-health problems resulting from high levels of Nr in water and land systems, and the human food supply (Vitousek et al., 1997; Britto and Kronzucker, 2002; Galloway et al., 2008). Moreover, while increases in anthropogenic N deposition have been increasing the size of terrestrial C sinks (Frey et al., 2014; Maaroufi et al., 2015), it is unlikely to keep pace with eCO_2 (Reay et al., 2008; Bala et al., 2013).




Fig. 3. Summary of the main aboveground and belowground responses to eCO_2 under low and high available N (see text for details).

3.5. Biological N₂ fixation (BNF)

Legumes and other N₂-fixing plants are able to overcome soil N limitations via symbiotic association with diazotrophic bacteria, and many studies have indicated that the extent of N₂ fixation increases under eCO₂ (Johnson, 2006; Reich et al., 2006b; Rogers et al., 2006; Kant et al., 2012), although not universally (Arnold, 1999; Hungate et al., 2004; van Groenigen et al., 2006). This has been observed in grasslands (Zanetti et al., 1996; Luscher et al., 2000), agricultural crops such as soybean (Torbert et al., 2004), and N₂-fixing trees such as alder (Koike et al., 2015). In addition, increased N₂ fixation has been observed among free-living cyanobacteria associated with rice roots in paddy soil (Hoque et al., 2001). Associated with this is a pronouncedly higher response in biomass gain in N₂-fixers relative to non-fixers (Rogers et al., 2006; cf. van Groenigen et al., 2006). In addition to the increase in N supply via nodulation, the commensal association of plant roots with diazotrophs provides an additional sink for growth stimulation under eCO₂, in the form of photosynthate transferred from plant to bacterium (Ainsworth and Rogers, 2007). The reduction in tissue N generally observed for C₃ plants is often not pronounced in legumes; for instance, Torbert et al. (2004) found a 31% increase in the C:N ratio in sorghum, but no change in this ratio in soybean.

Soil N supply can also constrain the response of BNF to eCO₂. It is generally viewed that N-fixation rates decrease with increased soil N availability (Vitousek et al., 2002; cf. Pearson and Vitousek, 2001). This seems to be the case as well under eCO₂; for example, in a pine forest, N deposition was shown to suppress BNF under eCO₂, suppressing the diversity and abundance of N₂-fixing bacteria (Berthrong et al., 2014; Fig. 3). Indeed, the stimulation in growth of legumes and rates of BNF under eCO₂ are often seen only under N-limiting conditions (Hungate et al., 1999; Poorter and Navas, 2003; West et al., 2005). However, as West et al. (2005) demonstrate, this response can vary significantly across species, with some legumes showing significant biomass gains with eCO₂ regardless of soil N supply (Lee et al., 2003). It is important to note, in addition, that even N₂-fixing plants can be strongly nutrient-limited, typically by the low availability of soil P, K, Mg, and Mo (Hungate et al., 2004; Reich et al., 2006a; van Groenigen et al., 2006; Rogers et al., 2009).

4. Conclusion

In this review, we have discussed the effects of eCO₂ and N limitation on photosynthesis, respiration, N transport and assimilation, C and N accumulation, and growth and yield in a variety of plant systems and ecosystems. In addition, we have examined effects on soil N processes, including PNL, N mineralization, and BNF. The study and integration of these plant and soil processes is essential not only to the understanding of vegetation and agronomic responses to eCO₂ and how they intersect with N supply, but also to help design strategies by which to mitigate CO₂ emissions resulting from human activities.

While it is clear that eCO₂ increases the NPP of most C₃ plants particularly strongly when N supply is adequate, however, the approach to artificially fertilize natural ecosystems with N to increase their C-sink capacity would be problematic. As mentioned, the synthesis of N fertilizer is energetically costly, and the amount of N₂ fixed in this manner is already approximately equal to the total N₂ fixed by Earth's diazotrophs, having brought about a major shift in the global N cycle, and entailing a host of problems for environmental and human health (Vitousek and Howarth, 1991; Fowler et al., 2015). Particularly worth noting are eutrophication effects (Vitousek and Howarth, 1991), as well as the finding that, while

eCO₂ can increase biodiversity (Reich, 2009), elevated N can lead to a loss of biodiversity (e.g. Langley and Megonigal, 2010; Isbell et al., 2013), which can negatively impact ecosystem functioning (Reich et al., 2001; Gamfeldt et al., 2013; Tilman et al., 2014). In one study, intermediate levels of species loss (21–40%) reduced primary production to levels comparable to that of climate warming, while higher levels of extirpation (41–60%) had effects comparable to ozone, acidification, eCO₂, and nutrient pollution (Hooper et al., 2012). Intermediate levels of species loss also had equal or greater effects on decomposition than eCO₂ or N deposition. In another study, decreases in grassland plant diversity influenced productivity at least as much as changes in nitrogen, water, CO₂, herbivores, drought, and fire (Tilman et al., 2012). Interestingly, in this study, the impact of biodiversity as a driver of ecosystem productivity increased over time, unlike the impact of other factors which either declined (i.e. nitrogen addition) or remained unchanged (all others).

In agricultural and silvicultural systems, where N is routinely supplemented, there may be potential for greenhouse-gas mitigation, as the growth and yield of crop species tend to respond well to eCO₂, especially so in the case of legumes (Jensen et al., 2012). Clearly, the intelligent management of soil resources will be required, such as the coordinated application of N at specific stages of plant development (Kim et al., 2001), and increases in the N-use efficiency of cropping systems, which currently stands at an average of only about 40% in the case of cereals (Kant et al., 2012). In addition, new attention will need to be focused on the source of N provided for fertilization, given the different responses of plants to NH₄⁺ and NO₃⁻, and possibly organic sources of N, under eCO₂ (Bauer and Berntson, 2001; Cruz et al., 2014; Bloom, 2015b). Secondly, advances in plant breeding to produce genotypes pre-adapted to changing environmental conditions will be very important in the coming decades. Target areas pertinent to the issues discussed here will include breeding plants with improved mineral nutrition, particularly that of N and P (Richardson et al., 2011; Pilbeam, 2015), and increased sink strength to maximize utilization of increasingly plant-available CO₂ (Wang et al., 2013). Advances in these areas specific to eCO₂ and N limitation will need to be coordinated with ongoing efforts to increase photosynthetic efficiency and productivity in plants to sustainably meet global food demands (Ort et al., 2015; White et al., 2016).

To conclude, we reiterate that accurate scientific models are critical to help policy makers design strategies to preserve, or increase, carbon sinks, and thus to mitigate eCO₂ and thereby climate change (Friedlingstein et al., 2011). Understanding the role of nutrient cycles in plant carbon fixation is essential to the development of reliable models (Zaehle et al., 2015), with a view to healthy natural-ecosystem function and the sustainable future of agricultural production, both of which are threatened by continued human population and resource-consumption growth and by the unprecedentedly rapid pace of global environmental change (Long et al., 2006; Lobell et al., 2011; Myers et al., 2014).

This review summarizes the effects of elevated atmospheric [CO₂] and N limitation on photosynthesis, respiration, N transport and assimilation, C and N accumulation, and growth and yield in a variety of plant systems and ecosystems. We highlight key studies from the field, particularly from free-air CO₂ enrichment (FACE) technology, and draw on both fundamental and novel physiological and ecological mechanisms for connection.

Acknowledgements

Funding was provided by the Natural Sciences and Engineering Research Council of Canada (NSERC).

References

- Adair, E.C., Reich, P.B., Trost, J.J., Hobbie, S.E., 2011. Elevated CO₂ stimulates grassland soil respiration by increasing carbon inputs rather than by enhancing soil moisture. *Glob. Change Biol.* 17, 3546–3563.
- Ainsworth, E.A., Long, S.P., 2005. What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy. *New Phytol.* 165, 351–371.
- Ainsworth, E.A., Rogers, A., 2007. The response of photosynthesis and stomatal conductance to rising CO₂: mechanisms and environmental interactions. *Plant Cell Environ.* 30, 258–270.
- Ainsworth, E.A., Davey, P.A., Bernacchi, C.J., Dermody, O.C., Heaton, E.A., Moore, D.J., Morgan, P.B., Naidu, S.L., Ra, H.S.Y., Zhu, X.G., Curtis, P.S., Long, S.P., 2002. A meta-analysis of elevated CO₂ effects on soybean (*Glycine max*) physiology, growth and yield. *Glob. Change Biol.* 8, 695–709.
- Ainsworth, E.A., Davey, P.A., Hymus, G.J., Osborne, C.P., Rogers, A., Blum, H., Nosberger, J., Long, S.P., 2003. Is stimulation of leaf photosynthesis by elevated carbon dioxide concentration maintained in the long term? A test with *Lolium perenne* grown for 10 years at two nitrogen fertilization levels under free air CO₂ enrichment (FACE). *Plant Cell Environ.* 26, 705–714.
- Andres, R.J., Boden, T.A., Breon, F.M., Ciais, P., Davis, S., Erickson, D., Gregg, J.S., Jacobson, A., Marland, G., Miller, J., Oda, T., Olivier, J.G.J., Raupach, M.R., Rayner, P., Treanton, K., 2012. A synthesis of carbon dioxide emissions from fossil-fuel combustion. *Biogeosciences* 9, 1845–1871.
- Arnth, A., Harrison, S.P., Zaehle, S., Tsigaridis, K., Menon, S., Bartlein, P.J., Feichter, J., Korhola, A., Kulmala, M., O'Donnell, D., Schurgers, G., Sorvari, S., Vesala, T., 2010. Terrestrial biogeochemical feedbacks in the climate system. *Nat. Geosci.* 3, 525–532.
- Arnold, J.A., 1999. Symbiotic N₂ fixation in a high Alpine grassland: effects of four growing seasons of elevated CO₂. *Funct. Ecol.* 13, 383–387.
- Arnoiz, P.A., Nelemans, J.A., Findenegg, G.R., 1988. Phosphoenolpyruvate carboxylase activity in plants grown with either NO₃⁻ or NH₄⁺ as inorganic nitrogen source. *J. Plant Physiol.* 132, 23–27.
- Arp, W.J., 1991. Effects of source-sink relations on photosynthetic acclimation to elevated CO₂. *Plant Cell Environ.* 14, 869–875.
- Bai, E., Li, S., Xu, W., Li, W., Dai, W., Jiang, P., 2013. A meta-analysis of experimental warming effects on terrestrial nitrogen pools and dynamics. *New Phytol.* 199, 441–451.
- Bala, G., Devaraju, N., Chaturvedi, R.K., Caldeira, K., Nemani, R., 2013. Nitrogen deposition: how important is it for global terrestrial carbon uptake? *Biogeosciences* 10, 7147–7160.
- Balkos, K.D., Britto, D.T., Kronzucker, H.J., 2010. Optimization of ammonium acquisition and metabolism by potassium in rice (*Oryza sativa* L. cv. IR-72). *Plant Cell Environ.* 33, 23–34.
- BassiriRad, H., Gutschick, V.P., Lussenhop, J., 2001. Root system adjustments: regulation of plant nutrient uptake and growth responses to elevated CO₂. *Oecologia* 126, 305–320.
- Bauer, G.A., Berntson, G.M., 2001. Ammonium and nitrate acquisition by plants in response to elevated CO₂ concentration: the roles of root physiology and architecture. *Tree Physiol.* 21, 137–144.
- Becker, C., Kläring, H.-P., 2016. CO₂ enrichment can produce high red leaf lettuce yield while increasing most flavonoid glycoside and some caffeic acid derivative concentrations. *Food Chem.* 199, 736–745.
- Beidler, K.V., Taylor, B.N., Strand, A.E., Cooper, E.R., Schoenholz, M., Pritchard, S.G., 2015. Changes in root architecture under elevated concentrations of CO₂ and nitrogen reflect alternate soil exploration strategies. *New Phytol.* 205, 1153–1163.
- Berthrong, S.T., Yeager, C.M., Gallegos-Graves, L., Steven, B., Eichorst, S.A., Jackson, R.B., Kuske, C.R., 2014. Nitrogen fertilization has a stronger effect on soil nitrogen-fixing bacterial communities than elevated atmospheric CO₂. *Appl. Environ. Microbiol.* 80, 3103–3112.
- Bloom, A.J., 2015a. Photorespiration and nitrate assimilation: a major intersection between plant carbon and nitrogen. *Photosynth. Res.* 123, 117–128.
- Bloom, A.J., 2015b. The increasing importance of distinguishing among plant nitrogen sources. *Curr. Opin. Plant Biol.* 25, 10–16.
- Bloom, A.J., Smart, D.R., Nguyen, D.T., Searles, P.S., 2002. Nitrogen assimilation and growth of wheat under elevated carbon dioxide. *Proc. Natl. Acad. Sci. U. S. A.* 99, 1730–1735.
- Bloom, A.J., Burger, M., Rubio-Asensio, J.S., Cousins, A.B., 2010. Carbon dioxide enrichment inhibits nitrate assimilation in wheat and *Arabidopsis*. *Science* 328, 899–903.
- Bowler, J.M., Press, M.C., 1996. Effects of elevated CO₂, nitrogen form and concentration on growth and photosynthesis of a fast- and slow-growing grass. *New Phytol.* 132, 391–401.
- Britto, D.T., Kronzucker, H.J., 2002. NH₄⁺ toxicity in higher plants: a critical review. *J. Plant Physiol.* 159, 567–584.
- Britto, D.T., Balkos, K.D., Becker, A., Coskun, D., Huynh, W.Q., Kronzucker, H.J., 2014. Potassium and nitrogen poisoning: physiological changes and biomass gains in rice and barley. *Can. J. Plant Sci.* 94, 1085–1089.
- Bunce, J.A., 2012. Responses of cotton and wheat photosynthesis and growth to cyclic variation in carbon dioxide concentration. *Photosynthetica* 50, 395–400.
- Bunce, J.A., 2014. CO₂ enrichment at night affects the growth and yield of common beans. *Crop Sci.* 54, 1744–1747.
- Burton, A.J., Pregitzer, K.S., Crawford, J.N., Zogg, G.P., Zak, D.R., 2004. Simulated chronic NO₃⁻ deposition reduces soil respiration in northern hardwood forests. *Glob. Change Biol.* 10, 1080–1091.
- Butler, S.M., Melillo, J.M., Johnson, J.E., Mohan, J., Steudler, P.A., Lux, H., Burrows, E., Smith, R.M., Vario, C.L., Scott, L., Hill, T.D., 2012. Soil warming alters nitrogen cycling in a New England forest: implications for ecosystem function and structure. *Oecologia* 168, 819–828.
- Cai, W., Borlace, S., Lengaigne, M., van Rensch, P., Collins, M., Vecchi, G., Timmermann, A., Santoso, A., McPhaden, M.J., Wu, L., England, M.H., Wang, G., Guilyardi, E., Jin, F.-F., 2014. Increasing frequency of extreme El Niño events due to greenhouse warming. *Nat. Clim. Change* 4, 111–116.
- Chang, J., Ciais, P., Viovy, N., Vuichard, N., Herrero, M., Havlík, P., Wang, X., Sultan, B., Soussana, J.F., 2016. Effect of climate change, CO₂ trends, nitrogen addition, and land-cover and management intensity changes on the carbon balance of European grasslands. *Glob. Change Biol.* 22, 338–350.
- Cheng, L., Booker, F.L., Tu, C., Burkey, K.O., Zhou, L., Shew, H.D., Rufty, T.W., Hu, S., 2012. Arbuscular mycorrhizal fungi increase organic carbon decomposition under elevated CO₂. *Science* 337, 1084–1087.
- Chollet, R., Vidal, J., Oleary, M.H., 1996. Phosphoenolpyruvate carboxylase: a ubiquitous, highly regulated enzyme in plants. *Ann. Rev. Plant Phys.* 47, 273–298.
- Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogee, J., Allard, V., Aubinet, M., Buchmann, N., Bernhofer, C., Carrara, A., Chevallier, F., De Noblet, N., Friend, A.D., Friedlingstein, P., Grunwald, T., Heinesch, B., Kerönen, P., Knohl, A., Krinner, G., Loustau, D., Manca, G., Matteucci, G., Miglietta, F., Ourcival, J.M., Papale, D., Pilegaard, K., Rambal, S., Seufert, G., Soussana, J.F., Sanz, M.J., Schulze, E.D., Vesala, T., Valentini, R., 2005. Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature* 437, 529–533.
- Cleveland, C.C., Townsend, A.R., Taylor, P., Alvarez-Clare, S., Bustamante, M.M.C., Chuyong, G., Dobrowski, S.Z., Grieron, P., Harms, K.E., Houlton, B.Z., Marklein, A., Parton, W., Porder, S., Reed, S.C., Sierra, C.A., Silver, W.L., Tanner, E.V.J., Wieder, W.R., 2011. Relationships among net primary productivity, nutrients and climate in tropical rain forest: a pan-tropical analysis. *Ecol. Lett.* 14, 939–947.
- Cleveland, C.C., Taylor, P., Chadwick, K.D., Dahlin, K., Doughty, C.E., Malhi, Y., Smith, W.K., Sullivan, B.W., Wieder, W.R., Townsend, A.R., 2015. A comparison of plot-based satellite and earth system model estimates of tropical forest net primary production. *Glob. Biogeochem. Cycles* 29, 626–644.
- Cortrufo, M.F., Ineson, P., Scott, A., 1998. Elevated CO₂ reduces the nitrogen concentration of plant tissues. *Glob. Change Biol.* 4, 43–54.
- Craine, J.M., Wedin, D.A., Reich, P.B., 2001. The response of soil CO₂ flux to changes in atmospheric CO₂, nitrogen supply and plant diversity. *Glob. Change Biol.* 7, 947–953.
- Cramer, M.D., Lewis, O.A.M., Lips, S.H., 1993. Inorganic carbon fixation and metabolism in maize roots as affected by nitrate and ammonium nutrition. *Physiol. Plant.* 89, 632–639.
- Cramer, W., Bondeau, A., Woodward, F.I., Prentice, I.C., Betts, R.A., Brovkin, V., Cox, P.M., Fisher, V., Foley, J.A., Friend, A.D., Kucharik, C., Lomas, M.R., Ramankutty, N., Sitch, S., Smith, B., White, A., Young-Molling, C., 2001. Global response of terrestrial ecosystem structure and function to CO₂ and climate change: results from six dynamic global vegetation models. *Glob. Change Biol.* 7, 357–373.
- Cramer, M.D., Hawkins, H.-J., Verboom, G.A., 2009. The importance of nutritional regulation of plant water flux. *Oecologia* 161, 15–24.
- Cruz, J.L., Alves, A.A.C., LeCain, D.R., Ellis, D.D., Morgan, J.A., 2014. Effect of elevated CO₂ concentration and nitrate:ammonium ratios on gas exchange and growth of cassava (*Manihot esculenta* Crantz). *Plant Soil* 374, 33–43.
- Curtis, P.S., Wang, X.Z., 1998. A meta-analysis of elevated CO₂ effects on woody plant mass, form, and physiology. *Oecologia* 113, 299–313.
- Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440, 165–173.
- de Graaff, M.-A., van Groenigen, K.-J., Six, J., Hungate, B., van Kessel, C., 2006. Interactions between plant growth and soil nutrient cycling under elevated CO₂: a meta-analysis. *Glob. Change Biol.* 12, 2077–2091.
- Dieleman, W.I.J., Vicca, S., Dijkstra, F.A., Hagedorn, F., Hovenden, M.J., Larsen, K.S., Morgan, J.A., Volder, A., Beier, C., Dukes, J.S., King, J., Leuzinger, S., Linder, S., Luo, Y.Q., Oren, R., de Angelis, P., Tingey, D., Hoosbeek, M.R., Janssens, I.A., 2012. Simple additive effects are rare: a quantitative review of plant biomass and soil process responses to combined manipulations of CO₂ and temperature. *Glob. Change Biol.* 18, 2681–2693.
- Dijkstra, P., Hymus, G., Colavito, D., Vieglais, D.A., Cundari, C.M., Johnson, D.P., Hungate, B.A., Hinkle, C.R., Drake, B.G., 2002. Elevated atmospheric CO₂ stimulates aboveground biomass in a fire-regenerated scrub-oak ecosystem. *Glob. Change Biol.* 8, 90–103.
- Dijkstra, F.A., Pendall, E., Mosier, A.R., King, J.Y., Milchunas, D.G., Morgan, J.A., 2008. Long-term enhancement of N availability and plant growth under elevated CO₂ in a semi-arid grassland. *Funct. Ecol.* 22, 975–982.
- Drake, B.G., Gonzalez-Meler, M.A., Long, S.P., 1997. More efficient plants: a consequence of rising atmospheric CO₂? *Ann. Rev. Plant Phys.* 48, 609–639.
- Drake, J.E., Gallet-Budynek, A., Hofmockel, K.S., Bernhardt, E.S., Billings, S.A., Jackson, R.B., Johnsen, K.S., Lichter, J., McCarthy, H.R., McCormack, M.L., Moore, D.J.P., Oren, R., Palmroth, S., Phillips, R.P., Pippin, J.S., Pritchard, S.G., Treseder, K.K., Schlesinger, W.H., DeLucia, E.H., Finzi, A.C., 2011. Increases in the flux of carbon belowground stimulate nitrogen uptake and sustain the long-term enhancement of forest productivity under elevated CO₂. *Ecol. Lett.* 14, 349–357.
- Dukes, J.S., Chiariello, N.R., Cleland, E.E., Moore, L.A., Shaw, M.R., Thayer, S., Tobeck, T., Mooney, H.A., Field, C.B., 2005. Responses of grassland production to single and multiple global environmental changes. *PLoS Biol.* 3, 1829–1837.

- Ellsworth, D.S., Reich, P.B., Naumburg, E.S., Koch, G.W., Kubiske, M.E., Smith, S.D., 2004. Photosynthesis, carboxylation and leaf nitrogen responses of 16 species to elevated pCO₂ across four free-air CO₂ enrichment experiments in forest, grassland and desert. *Glob. Change Biol.* 10, 2121–2138.
- Evans, J.R., 1989. Photosynthesis and nitrogen relationships in leaves of C₃ plants. *Oecologia* 78, 9–19.
- Eviner, V.T., Chapin, F.S., 1997. Nitrogen cycle—plant-microbial interactions. *Nature* 385, 26–27.
- Fangmeier, A., De Temmerman, L., Mortensen, L., Kemp, K., Burke, J., Mitchell, R., van Oijen, M., Weigel, H.J., 1999. Effects on nutrients and on grain quality in spring wheat crops grown under elevated CO₂ concentrations and stress conditions in the European, multiple-site experiment 'ESPACE-wheat'. *Eur. J. Agron.* 10, 215–229.
- Farage, P.K., McKee, I.F., Long, S.P., 1998. Does a low nitrogen supply necessarily lead to acclimation of photosynthesis to elevated CO₂? *Plant Physiol.* 118, 573–580.
- Feng, Z., Ruetting, T., Pleijel, H., Wallin, G., Reich, P.B., Kammann, C.I., Newton, P.C.D., Kobayashi, K., Luo, Y., Uddling, J., 2015. Constraints to nitrogen acquisition of terrestrial plants under elevated CO₂. *Glob. Change Biol.* 21, 3152–3168.
- Fernández, M.D., Tezara, W., Rengifo, E., Herrera, A., 2002. Lack of downregulation of photosynthesis in a tropical root crop, cassava, grown under an elevated CO₂ concentration. *Funct. Plant Biol.* 29, 805–814.
- Fernández-Martínez, M., Vicca, S., Janssens, I.A., Sardans, J., Luysaert, S., Campioli, M., Chapin, F.S., Ciais, P., Malhi, Y., Obersteiner, M., Papale, D., Piao, S.L., Reichstein, M., Roda, F., Penuelas, J., 2014. Nutrient availability as the key regulator of global forest carbon balance. *Nat. Clim. Change* 4, 471–476.
- Finzi, A.C., Schlesinger, W.H., 2003. Soil-nitrogen cycling in a pine forest exposed to 5 years of elevated carbon dioxide. *Ecosystems* 6, 444–456.
- Finzi, A.C., Norby, R.J., Calapietra, C., Gallet-Budynek, A., Gielen, B., Holmes, W.E., Hoosbeek, M.R., Iversen, C.M., Jackson, R.B., Kubiske, M.E., Ledford, J., Liberloo, M., Oren, R., Polle, A., Pritchard, S., Zak, D.R., Schlesinger, W.H., Ceulemans, R., 2007. Increases in nitrogen uptake rather than nitrogen-use efficiency support higher rates of temperate forest productivity under elevated CO₂. *Proc. Natl. Acad. Sci. U. S. A.* 104, 14014–14019.
- Forkel, M., Carvalhais, N., Rödenbeck, C., Keeling, R., Heimann, M., Thonicke, K., Zaehle, S., Reichstein, M., 2016. Enhanced seasonal CO₂ exchange caused by amplified plant productivity in northern ecosystems. *Science*, <http://dx.doi.org/10.1126/science.aac4971>.
- Fowler, D., Steadman, C.E., Stevenson, D., Coyle, M., Rees, R.M., Skiba, U.M., Sutton, M.A., Cape, J.N., Dore, A.J., Viero, M., Simpson, D., Zaehle, S., Stocker, B.D., Rinaldi, M., Fächini, M.C., Flechard, C.R., Nemitz, E., Twigg, M., Erisman, J.W., Butterbach-Bahl, K., Galloway, J.N., 2015. Effects of global change during the 21st century on the nitrogen cycle. *Atmos. Chem. Phys.* 15, 13849–13893.
- Frölicher, T.L., Winton, M., Sarmiento, J.L., 2014. Continued global warming after CO₂ emissions stoppage. *Nat. Clim. Change* 4, 40–44.
- Franks, P.J., Adams, M.A., Amthor, J.S., Barbour, M.M., Berry, J.A., Ellsworth, D.S., Farquhar, G.D., Ghanoum, O., Lloyd, J., McDowell, N., Norby, R.J., Tissue, D.T., von Caemmerer, S., 2013. Sensitivity of plants to changing atmospheric CO₂ concentration: from the geological past to the next century. *New Phytol.* 197, 1077–1094.
- Frey, S.D., Ollinger, S., Nadelhoffer, K., Bowden, R., Brzostek, E., Burton, A., Caldwell, B.A., Crow, S., Goodale, C.L., Grandy, A.S., Finzi, A.C., Kramer, M.G., Lajtha, K., LeMoine, J., Martin, M., McDowell, W.H., Minocha, R., Sadowsky, J.J., Templer, P.H., Wickings, K., 2014. Chronic nitrogen additions suppress decomposition and sequester soil carbon in temperate forests. *Biogeochemistry* 121, 305–316.
- Friedlingstein, P., Solomon, S., Plattner, G.K., Knutti, R., Ciais, P., Raupach, M.R., 2011. Long-term climate implications of twenty-first century options for carbon dioxide emission mitigation. *Nat. Clim. Change* 1, 457–461.
- Galloway, J.N., Dentener, F.J., Capone, D.G., Boyer, E.W., Howarth, R.W., Seitzinger, S.P., Asner, G.P., Cleveland, C.C., Green, P.A., Holland, E.A., Karl, D.M., Michaels, A.F., Porter, J.H., Townsend, A.R., Vorosmarty, C.J., 2004. Nitrogen cycles: past, present, and future. *Biogeochemistry* 70, 153–226.
- Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z., Freney, J.R., Martinelli, L.A., Seitzinger, S.P., Sutton, M.A., 2008. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science* 320, 889–892.
- Gamfeldt, L., Snäll, T., Bagchi, R., Jonsson, M., Gustafsson, L., Kjellander, P., Ruiz-Jaen, M.C., Fröberg, M., Stendahl, J., Philipson, C.D., Mikusiński, G., 2013. Higher levels of multiple ecosystem services are found in forests with more tree species. *Nat. Commun.* 4, 1340.
- Geiger, M., Walch-Liu, P., Engels, C., Harnecker, J., Schulze, E.D., Ludewig, F., Sonnewald, U., Scheible, W.R., Stitt, M., 1998. Enhanced carbon dioxide leads to a modified diurnal rhythm of nitrate reductase activity in older plants, and a large stimulation of nitrate reductase activity and higher levels of amino acids in young tobacco plants. *Plant Cell Environ.* 21, 253–268.
- Geiger, M., Haake, V., Ludewig, F., Sonnewald, U., Stitt, M., 1999. The nitrate and ammonium nitrate supply have a major influence on the response of photosynthesis carbon metabolism, nitrogen metabolism and growth to elevated carbon dioxide in tobacco. *Plant Cell Environ.* 22, 1177–1199.
- Gifford, R.M., Barrett, D.J., Lütze, J.L., 2000. The effects of elevated CO₂ on the C:N and C:P mass ratios of plant tissues. *Plant Soil* 224, 1–14.
- Gigon, A., Rorison, I.H., 1972. Response of some ecologically distinct plant species to nitrate-nitrogen and to ammonium-nitrogen. *J. Ecol.* 60, 93–102.
- Gill, R.A., Polley, H.W., Johnson, H.B., Anderson, L.J., Maherali, H., Jackson, R.B., 2002. Nonlinear grassland responses to past and future atmospheric CO₂. *Nature* 417, 279–282.
- Givan, C.V., 1979. Metabolic detoxification of ammonia in tissues of higher plants. *Phytochemistry* 18, 375–382.
- Guo, S., Zhou, Y., Shen, Q., Zhang, F., 2007. Effect of ammonium and nitrate nutrition on some physiological processes in higher plants—growth photosynthesis, photorespiration, and water relations. *Plant Biol.* 9, 21–29.
- Guo, H., Sun, Y., Li, Y., Liu, X., Ren, Q., Zhu-Salzman, K., Ge, F., 2013. Elevated CO₂ modifies N acquisition of *Medicago truncatula* by enhancing N fixation and reducing nitrate uptake from soil. *PLoS One* 8, e81373.
- Högy, P., Wieser, H., Köhler, P., Schwadorf, K., Breuer, J., Erbs, M., Weber, S., Fangmeier, A., 2009. Does elevated atmospheric CO₂ allow for sufficient wheat grain quality in the future? *J. Appl. Bot. Food Qual.* 82, 114–121.
- Harley, P.C., Thomas, R.B., Reynolds, J.F., Strain, B.R., 1992. Modeling photosynthesis of cotton grown in elevated CO₂. *Plant Cell Environ.* 15, 271–282.
- Harvell, C.D., Mitchell, C.E., Ward, J.R., Altizer, S., Dobson, A.P., Ostfeld, R.S., Samuel, M.D., 2002. Ecology—climate warming and disease risks for terrestrial and marine biota. *Science* 296, 2158–2162.
- Hasselquist, N.J., Metcalfe, D.B., Hogberg, P., 2012. Contrasting effects of low and high nitrogen additions on soil CO₂ flux components and ectomycorrhizal fungal sporocarp production in a boreal forest. *Glob. Change Biol.* 18, 3596–3605.
- Haynes, R.J., Goh, K.M., 1978. Ammonium and nitrate nutrition of plants. *Biol. Rev. Camb. Philos.* 53, 465–510.
- Hofmockel, K.S., Schlesinger, W.H., Jackson, R.B., 2007. Effects of elevated atmospheric carbon dioxide on amino acid and NH₄⁺-N cycling in a temperate pine ecosystem. *Glob. Change Biol.* 13, 1950–1959.
- Hofmockel, K.S., Fierer, N., Colman, B.P., Jackson, R.B., 2010. Amino acid abundance and proteolytic potential in North American soils. *Oecologia* 163, 1069–1078.
- Hofmockel, K.S., Gallet-Budynek, A., McCarthy, H.R., Currie, W.S., Jackson, R.B., Finzi, A.C., 2011. Sources of increased N uptake in forest trees growing under elevated CO₂: results of a large-scale N-15 study. *Glob. Change Biol.* 17, 3338–3350.
- Hooper, D.U., Adair, E.C., Cardinale, B.J., Byrnes, J.E., Hungate, B.A., Matulich, K.L., Gonzalez, A., Duffy, J.E., Gamfeldt, L., O'Connor, M.I., 2012. A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* 486, 105–108.
- Hoque, M.M., Inubushi, K., Miura, S., Kobayashi, K., Kim, H.Y., Okada, M., Yabashi, S., 2001. Biological dinitrogen fixation and soil microbial biomass carbon as influenced by free-air carbon dioxide enrichment (FACE) at three levels of nitrogen fertilization in a paddy field. *Biol. Fertil. Soils* 34, 453–459.
- Hovenden, M.J., Newton, P.C.D., Carran, R.A., Theobald, P., Wills, K.E., Schoor, J.K.V., Williams, A.L., Osanai, Y., 2008. Warming prevents the elevated CO₂-induced reduction in available soil nitrogen in a temperate, perennial grassland. *Glob. Change Biol.* 14, 1018–1024.
- Hungate, B.A., Dijkstra, P., Johnson, D., Hinkle, C.R., Drake, B., 1999. Elevated CO₂ increases nitrogen fixation and decreases soil nitrogen mineralization in Florida scrub oak. *Glob. Change Biol.* 5, 781–789.
- Hungate, B.A., Dukes, J.S., Shaw, M.R., Luo, Y.Q., Field, C.B., 2003. Nitrogen and climate change. *Science* 302, 1512–1513.
- Hungate, B.A., Stiling, P.D., Dijkstra, P., Johnson, D.W., Ketterer, M.E., Hymus, G.J., Hinkle, C.R., Drake, B.G., 2004. CO₂ elicits long-term decline in nitrogen fixation. *Science* 304, 1291.
- Hungate, B.A., Johnson, D.W., Dijkstra, P., Hymus, G., Stiling, P., Megonigal, J.P., Pagel, A.L., Moan, J.L., Day, F., Li, J.H., Hinkle, C.R., Drake, B.G., 2006. Nitrogen cycling during seven years of atmospheric CO₂ enrichment in a scrub oak woodland. *Ecology* 87, 26–40.
- Huppe, H.C., Turpin, D.H., 1994. Integration of carbon and nitrogen metabolism in plant and algal cells. *Ann. Rev. Plant Phys.* 45, 577–607.
- Hyvönen, R., Ågren, G.I., Linder, S., Persson, T., Cotrufo, M.F., Ekblad, A., Freeman, M., Grelle, A., Janssens, I.A., Jarvis, P.G., Kellomäki, S., Lindroth, A., Loustau, D., Lundmark, T., Norby, R.J., Oren, R., Pilegaard, K., Ryan, M.G., Sigurdsson, B.D., Strömgren, M., van Oijen, M., Wallin, G., 2007. The likely impact of elevated CO₂, nitrogen deposition, increased temperature and management on carbon sequestration in temperate and boreal forest ecosystems: a literature review. *New Phytol.* 173, 463–480.
- Ikedo, M., Mizoguchi, K., Yamakawa, T., 1992. Stimulation of dark carbon fixation in rice and tomato roots by application of ammonium nitrogen. *Soil Sci. Plant Nutr.* 38, 315–322.
- Imsande, J., Touraine, B., 1994. N demand and the regulation of nitrate uptake. *Plant Physiol.* 105, 3–7.
- Inauen, N., Körner, C., Hiltbrunner, E., 2012. No growth stimulation by CO₂ enrichment in alpine glacier forefield plants. *Glob. Change Biol.* 18, 985–999.
- Isbell, F., Reich, P.B., Tilman, D., Hobbie, S.E., Polasky, S., Binder, S., 2013. Nutrient enrichment, biodiversity loss, and consequent declines in ecosystem productivity. *Proc. Natl. Acad. Sci. U. S. A.* 110, 11911–11916.
- Iversen, C.M., Hooker, T.D., Classen, A.T., Norby, R.J., 2011. Net mineralization of N at deeper soil depths as a potential mechanism for sustained forest production under elevated CO₂. *Glob. Change Biol.* 17, 1130–1139.
- Iversen, C.M., Keller, J.K., Garten, C.T., Norby, R.J., 2012. Soil carbon and nitrogen cycling and storage throughout the soil profile in a sweetgum plantation after 11 years of CO₂-enrichment. *Glob. Change Biol.* 18, 1684–1697.
- Jablonski, L.M., Wang, X.Z., Curtis, P.S., 2002. Plant reproduction under elevated CO₂ conditions: a meta-analysis of reports on 79 crop and wild species. *New Phytol.* 156, 9–26.

- Jackson, L.E., Burger, M., Cavignaro, T.R., 2008. **Roots nitrogen transformations, and ecosystem services.** *Ann. Rev. Plant Biol.* 59, 341–363.
- Janssens, I.A., Dieleman, W., Luysaert, S., Subke, J.A., Reichstein, M., Ceulemans, R., Ciais, P., Dolman, A.J., Grace, J., Matteucci, G., Papale, D., Piao, S.L., Schulze, E.D., Tang, J., Law, B.E., 2010. **Reduction of forest soil respiration in response to nitrogen deposition.** *Nat. Geosci.* 3, 315–322.
- Jauregui, I., Aparicio-Tejo, P.M., Avila, C., Cañas, R., Sakalauskiene, S., Aranjuelo, I., 2016. **Root-shoot interactions explain the reduction of leaf mineral content in *Arabidopsis* plants grown under elevated $[CO_2]$ conditions.** *Physiol. Plant.*, <http://dx.doi.org/10.1111/pp1.12417>.
- Jensen, R.G., 2000. **Activation of Rubisco regulates photosynthesis at high temperature and CO_2 .** *Proc. Natl. Acad. Sci. U. S. A.* 97, 12937–12938.
- Jensen, E.S., Peoples, M.B., Boddey, R.M., Gresshoff, P.M., Hauggaard-Nielsen, H., Alves, B.J.R., Morrison, M.J., 2012. **Legumes for mitigation of climate change and the provision of feedstock for biofuels and biorefineries: a review.** *Agron. Sustain. Dev.* 32, 329–364.
- Johnson, D.W., 2006. **Progressive N limitation in forests: review and implications for long-term responses to elevated CO_2 .** *Ecology* 87, 64–75.
- Jones, A.G., Scullion, J., Ostle, N., Levy, P.E., Gwynn-Jones, D., 2014. **Completing the FACE of elevated CO_2 research.** *Environ. Int.* 73, 252–258.
- Körner, C., 2006. **Plant CO_2 responses: an issue of definition, time and resource supply.** *New Phytol.* 172, 393–411.
- Körner, C., 2015. **Paradigm shift in plant growth control.** *Curr. Opin. Plant Biol.* 25, 107–114.
- Kaiser, W.M., Forster, J., 1989. **Low CO_2 prevents nitrate reduction in leaves.** *Plant Physiol.* 91, 970–974.
- Kant, S., Seneweera, S., Rodin, J., Materne, M., Burch, D., Rothstein, S.J., Spangenberg, G., 2012. **Improving yield potential in crops under elevated CO_2 : integrating the photosynthetic and nitrogen utilization efficiencies.** *Front. Plant Sci.* 3, 1–9.
- Kaye, J.P., Hart, S.C., 1997. **Competition for nitrogen between plants and soil microorganisms.** *Trends Ecol. Evol.* 12, 139–143.
- Kim, H.Y., Lieffering, M., Miura, S., Kobayashi, K., Okada, M., 2001. **Growth and nitrogen uptake of CO_2 -enriched rice under field conditions.** *New Phytol.* 150, 223–229.
- Kim, H.Y., Lieffering, M., Kobayashi, K., Okada, M., Miura, S., 2003. **Seasonal changes in the effects of elevated CO_2 on rice at three levels of nitrogen supply: a free air CO_2 enrichment (FACE) experiment.** *Glob. Change Biol.* 9, 826–837.
- Kimball, B.A., Kobayashi, K., Bindi, M., 2002. **Responses of agricultural crops to free-air CO_2 enrichment.** *Adv. Agron.* 77, 293–368.
- Kirk, G.J.K., Kronzucker, H.J., 2005. **The potential for nitrification and nitrate uptake in the rhizospheres of wetland plants: a modelling study.** *Ann. Bot. London* 96, 639–646.
- Klingensmith, K.M., Van Cleve, K., 1993. **Patterns of nitrogen mineralization and nitrification in floodplain successional soils along the Tanana river, interior Alaska.** *Can. J. For. Res.* 23, 964–969.
- Koga, N., Ikeda, M., 2000. **Methionine sulfoximine suppressed the stimulation of dark carbon fixation by ammonium nutrition in wheat roots.** *Soil Sci. Plant Nutr.* 46, 393–400.
- Koike, T., Watanabe, M., Watanabe, Y., Agathokleous, E., Eguchi, N., Takagi, K., Satoh, F., Kitaoka, S., Funada, R., 2015. **Ecophysiology of deciduous trees native to Northeast Asia grown under FACE (free air CO_2 enrichment).** *J. Agrometeorol.* 71, 174–184.
- Kronzucker, H.J., Siddiqi, M.Y., Glass, A.D.M., 1997. **Conifer root discrimination against soil nitrate and the ecology of forest succession.** *Nature* 385, 59–61.
- Langley, J.A., Megonigal, J.P., 2010. **Ecosystem response to elevated CO_2 levels limited by nitrogen-induced plant species shift.** *Nature* 466, 96–99.
- Langley, J.A., McKinley, D.C., Wolf, A.A., Hungate, B.A., Drake, B.G., Megonigal, J.P., 2009. **Priming depletes soil carbon and releases nitrogen in a scrub-oak ecosystem exposed to elevated CO_2 .** *Soil Biol. Biochem.* 41, 54–60.
- Larigauderie, A., Hilbert, D.W., Oechel, W.C., 1988. **Effect of CO_2 enrichment and nitrogen availability on resource acquisition and resource allocation in a grass, *Bromus mollis*.** *Oecologia* 77, 544–549.
- Le Quéré, C., Moriarty, R., Andrew, R.M., Canadell, J.G., Sitch, S., Korsbakken, J.I., Friedlingstein, P., Peters, G.P., Andres, R.J., Boden, T.A., Houghton, R.A., House, J.I., Keeling, R.F., Tans, P., Arneeth, A., Bakker, D.C.E., Barbero, L., Bopp, L., Chang, J., Chevallier, F., Chini, L.P., Ciais, P., Fader, M., Feely, R.A., Gkritzalis, T., Harris, I., Hauck, J., Ilyina, T., Jain, A.K., Kato, E., Kitidis, V., Klein Goldewijk, K., Koven, C., Landschützer, P., Lauvset, S.K., Lefèvre, N., Lenton, A., Lima, I.D., Metzl, N., Millero, F., Munro, D.R., Murata, A., Nabel, J.E.M.S., Nakaoka, S., Nojiri, Y., O'Brien, K., Olsen, A., Ono, T., Pérez, F.F., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Rödenbeck, C., Saito, S., Schuster, U., Schwinger, J., Séférian, R., Steinhoff, T., Stocker, B.D., Sutton, A.J., Takahashi, T., Tilbrook, B., van der Laan-Luijkx, I.T., van der Werf, G.R., van Heuven, S., Vandemark, D., Viovy, N., Wiltshire, A., Zaehle, S., Zeng, N., 2015. **Global carbon budget 2015.** *Earth Syst. Sci. Data* 7, 349–396.
- LeBauer, D.S., Treseder, K.K., 2008. **Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed.** *Ecology* 89, 371–379.
- Leadley, P.W., Drake, B.G., 1993. **Open top chambers for exposing plant canopies to elevated CO_2 concentration and for measuring net gas exchange.** *Vegetatio* 104, 3–15.
- Leakey, A.D.B., Ainsworth, E.A., Bernacchi, C.J., Rogers, A., Long, S.P., Ort, D.R., 2009. **Elevated CO_2 effects on plant carbon, nitrogen, and water relations: six important lessons from FACE.** *J. Exp. Bot.* 60, 2859–2876.
- Leakey, A.D.B., Bishop, K.A., Ainsworth, E.A., 2012. **A multi-biome gap in understanding of crop and ecosystem responses to elevated CO_2 .** *Curr. Opin. Plant Biol.* 15, 228–236.
- Lee, T.D., Tjoelker, M.G., Ellsworth, D.S., Reich, P.B., 2001. **Leaf gas exchange responses of 13 prairie grassland species to elevated CO_2 and increased nitrogen supply.** *New Phytol.* 150, 405–418.
- Lee, T.D., Tjoelker, M.G., Reich, P.B., Russelle, M.P., 2003. **Contrasting growth response of an N2-fixing and non-fixing forb to elevated CO_2 : dependence on soil N supply.** *Plant Soil* 255, 475–486.
- Lee, T.D., Barrott, S.H., Reich, P.B., 2011. **Photosynthetic responses of 13 grassland species across 11 years of free-air CO_2 enrichment is modest, consistent and independent of N supply.** *Glob. Change Biol.* 17, 2893–2904.
- Lenka, N.K., Lal, R., 2012. **Soil-related constraints to the carbon dioxide fertilization effect.** *Crit. Rev. Plant Sci.* 31, 342–357.
- Liu, H., Yang, L., Wang, Y., Huang, J., Zhu, J., Wang, Y., Dong, G., Liu, G., 2008. **Yield formation of CO_2 -enriched hybrid rice cultivar Shanyou 63 under fully open-air field conditions.** *Field Crops Res.* 108, 93–103.
- Lobell, D.B., Schlenker, W., Costa-Roberts, J., 2011. **Climate trends and global crop production since 1980.** *Science* 333, 616–620.
- Long, S.P., Ainsworth, E.A., Rogers, A., Ort, D.R., 2004. **Rising atmospheric carbon dioxide: plants face the future.** *Ann. Rev. Plant Biol.* 55, 591–628.
- Long, S.P., Ainsworth, E.A., Leakey, A.D.B., Nosberger, J., Ort, D.R., 2006. **Food for thought: lower-than-expected crop yield stimulation with rising CO_2 concentrations.** *Science* 312, 1918–1921.
- Lopes, M.S., Araus, J.L., van Heerden, P.D.R., Foyer, C.H., 2011. **Enhancing drought tolerance in C_4 crops.** *J. Exp. Bot.* 62, 3135–3153.
- Lucht, W., Prentice, I.C., Myneni, R.B., Sitch, S., Friedlingstein, P., Cramer, W., Bousquet, P., Buermann, W., Smith, B., 2002. **Climatic control of the high-latitude vegetation greening trend and Pinatubo effect.** *Science* 296, 1687–1689.
- Luo, Y., Field, C.B., Mooney, H.A., 1994. **Predicting responses of photosynthesis and root fraction to elevated CO_2 —interactions among carbon nitrogen, and growth.** *Plant Cell Environ.* 17, 1195–1204.
- Luo, Y.Q., Jackson, R.B., Field, C.B., Mooney, H.A., 1996. **Elevated CO_2 increases belowground respiration in California grasslands.** *Oecologia* 108, 130–137.
- Luo, Y., Su, B., Currie, W.S., Dukes, J.S., Finzi, A.C., Hartwig, U., Hungate, B., McMurtrie, R.E., Oren, R., Parton, W.J., Pataki, D.E., Shaw, M.R., Zak, D.R., Field, C.B., 2004. **Progressive nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide.** *Bioscience* 54, 731–739.
- Luo, Y.Q., Hui, D.F., Zhang, D.Q., 2006. **Elevated CO_2 stimulates net accumulations of carbon and nitrogen in land ecosystems: a meta-analysis.** *Ecology* 87, 53–63.
- Luscher, A., Hartwig, U.A., Suter, D., Nosberger, J., 2000. **Direct evidence that symbiotic N_2 fixation in fertile grassland is an important trait for a strong response of plants to elevated atmospheric CO_2 .** *Glob. Change Biol.* 6, 655–662.
- Ma, H., Zhu, J., Xie, Z., Liu, G., Zeng, Q., Han, Y., 2007. **Responses of rice and winter wheat to free-air CO_2 enrichment (China FACE) at rice/wheat rotation system.** *Plant Soil* 294, 137–146.
- Maaroufi, N.I., Nordin, A., Hasselquist, N.J., Bach, L.H., Palmqvist, K., Gundale, M.J., 2015. **Anthropogenic nitrogen deposition enhances carbon sequestration in boreal soils.** *Glob. Change Biol.* 21, 3169–3180.
- Magnani, F., Mencuccini, M., Borghetti, M., Berbigier, P., Berninger, F., Delzon, S., Grelle, A., Hari, P., Jarvis, P.G., Kolari, P., Kowalski, A.S., Lankreijer, H., Law, B.E., Lindroth, A., Loustau, D., Manca, G., Moncrieff, J.B., Rayment, M., Tedeschi, V., Valentini, R., Grace, J., 2007. **The human footprint in the carbon cycle of temperate and boreal forests.** *Nature* 447, 848–850.
- Matimati, I., Verboom, G.A., Cramer, M.D., 2014. **Nitrogen regulation of transpiration controls mass-flow acquisition of nutrients.** *J. Exp. Bot.* 65, 159–168.
- Matt, P., Geiger, M., Walch-Liu, P., Engels, C., Krapp, A., Stitt, M., 2001a. **The immediate cause of the diurnal changes of nitrogen metabolism in leaves of nitrate-replete tobacco: a major imbalance between the rate of nitrate reduction and the rates of nitrate uptake and ammonium metabolism during the first part of the light period.** *Plant Cell Environ.* 24, 177–190.
- Matt, P., Geiger, M., Walch-Liu, P., Engels, C., Krapp, A., Stitt, M., 2001b. **Elevated carbon dioxide increases nitrate uptake and nitrate reductase activity when tobacco is growing on nitrate, but increases ammonium uptake and inhibits nitrate reductase activity when tobacco is growing on ammonium nitrate.** *Plant Cell Environ.* 24, 1119–1137.
- Matthews, H.D., 2007. **Implications of CO_2 fertilization for future climate change in a coupled climate-carbon model.** *Glob. Change Biol.* 13, 1068–1078.
- McCarthy, H.R., Oren, R., Johnsen, K.H., Gallet-Budnyek, A., Pritchard, S.G., Cook, C.W., LaDeau, S.L., Jackson, R.B., Finzi, A.C., 2010. **Re-assessment of plant carbon dynamics at the Duke free-air CO_2 enrichment site: interactions of atmospheric CO_2 with nitrogen and water availability over stand development.** *New Phytol.* 185, 514–528.
- McDonald, E.P., Erickson, J.E., Kruger, E.L., 2002. **Can decreased transpiration limit plant nitrogen acquisition in elevated CO_2 ?** *Funct. Plant Biol.* 29, 1115–1120.
- McGrath, J.M., Lobell, D.B., 2013. **Reduction of transpiration and altered nutrient allocation contribute to nutrient decline of crops grown in elevated CO_2 concentrations.** *Plant Cell Environ.* 36, 697–705.
- McLeod, A.R., Long, S.P., 1999. **Free-air carbon dioxide enrichment (FACE) in global change research: a review.** *Adv. Ecol. Res.* 28, 1–56.
- McMurtrie, R.E., Norby, R.J., Medlyn, B.E., Dewar, R.C., Pepper, D.A., Reich, P.B., Barton, C.V.M., 2008. **Why is plant-growth response to elevated CO_2 amplified when water is limiting, but reduced when nitrogen is limiting? A growth-optimisation hypothesis.** *Funct. Plant Biol.* 35, 521–534.
- Medlyn, B.E., Badeck, F.W., De Pury, D.G.G., Barton, C.V.M., Broadmeadow, M., Ceulemans, R., De Angelis, P., Forstreuter, M., Jach, M.E., Kellomaki, S., Laitat, E., Marek, M., Philippot, S., Rey, A., Strassmeyer, J., Laitinen, K., Lionzo, R., Portier,

- B., Roberntz, P., Wang, K., Jarvis, P.G., 1999. Effects of elevated CO₂ on photosynthesis in European forest species: a meta-analysis of model parameters. *Plant Cell Environ.* 22, 1475–1495.
- Melillo, J.M., McGuire, A.D., Kicklighter, D.W., Moore, B., Vorosmarty, C.J., Schloss, A.L., 1993. Global climate change and terrestrial net primary production. *Nature* 363, 234–240.
- Melillo, J.M., Butler, S., Johnson, J., Mohan, J., Steudler, P., Lux, H., Burrows, E., Bowles, F., Smith, R., Scott, L., Vario, C., Hill, T., Burton, A., Zhou, Y.-M., Tang, J., 2011. Oil warming, carbon-nitrogen interactions, and forest carbon budgets. *Proc. Natl. Acad. Sci. U. S. A.* 108, 9508–9512.
- Messerli, J., Bertrand, A., Bourassa, J., Belanger, G., Castonguay, Y., Tremblay, G., Baron, V., Seguin, P., 2015. Performance of low-cost open-top chambers to study long-term effects of carbon dioxide and climate under field conditions. *Agron. J.* 107, 916–920.
- Meyerholt, J., Zaehle, S., 2015. The role of stoichiometric flexibility in modelling forest ecosystem responses to nitrogen fertilization. *New Phytol.* 208, 1042–1055.
- Miglietta, F., Giuntoli, A., Bindi, M., 1996. The effect of free air carbon dioxide enrichment (FACE) and soil nitrogen availability on the photosynthetic capacity of wheat. *Photosynth. Res.* 47, 281–290.
- Moore, B.D., Cheng, S.H., Sims, D., Seemann, J.R., 1999. The biochemical and molecular basis for photosynthetic acclimation to elevated atmospheric CO₂. *Plant Cell Environ.* 22, 567–582.
- Mueller, K.E., Hobbie, S.E., Tilman, D., Reich, P.B., 2013. Effects of plant diversity, N fertilization, and elevated carbon dioxide on grassland soil N cycling in a long-term experiment. *Glob. Change Biol.* 19, 1249–1261.
- Myers, S.S., Zanolletti, A., Kloog, I., Huybers, P., Leakey, A.D.B., Bloom, A.J., Carlisle, E., Dietterich, L.H., Fitzgerald, G., Hasegawa, T., Holbrook, N.M., Nelson, R.L., Ottman, M.J., Raboy, V., Sakai, H., Sartor, K.A., Schwartz, J., Seneweera, S., Tausz, M., Usui, Y., 2014. Increasing CO₂ threatens human nutrition. *Nature* 510, 139–142.
- Myneni, R.B., Keeling, C.D., Tucker, C.J., Asrar, G., Nemani, R.R., 1997. Increased plant growth in the northern high latitudes from 1981 to 1991. *Nature* 386, 698–702.
- Mystakidis, S., Davin, E.L., Gruber, N., Seneviratne, S.I., 2016. Constraining future terrestrial carbon cycle projections using observation-based water and carbon flux estimates. *Glob. Change Biol.*, <http://dx.doi.org/10.1111/gcb.13217>.
- Norby, R.J., Zak, D.R., 2011. Ecological lessons from free-air CO₂ enrichment (FACE) experiments. *Ann. Rev. Ecol. Syst.* 42, 181–203.
- Norby, R.J., DeLucia, E.H., Gielen, B., Calafapietra, C., Giardina, C.P., King, J.S., Ledford, J., McCarthy, H.R., Moore, D.J.P., Ceulemans, R., De Angelis, P., Finzi, A.C., Karnosky, D.F., Kubiske, M.E., Lukac, M., Pregitzer, K.S., Scarascia-Mugnozza, G.E., Schlesinger, W.H., Oren, R., 2005. Forest response to elevated CO₂ is conserved across a broad range of productivity. *Proc. Natl. Acad. Sci. U. S. A.* 102, 18052–18056.
- Norby, R.J., Warren, J.M., Iversen, C.M., Medlyn, B.E., McMurtrie, R.E., 2010. CO₂ enhancement of forest productivity constrained by limited nitrogen availability. *Proc. Natl. Acad. Sci. U. S. A.* 107, 19368–19373.
- Norby, R.J., De Kauwe, M.G., Domingues, T.F., Duursma, R.A., Ellsworth, D.S., Goll, D.S., Lapola, D.M., Luus, K.A., MacKenzie, A.R., Medlyn, B.E., Pavlick, R., Rammig, A., Smith, B., Thomas, R., Thonicke, K., Walker, A.P., Yang, X., Zaehle, S., 2016. Model-data synthesis for the next generation of forest free-air CO₂ enrichment (FACE) experiments. *New Phytol.* 209, 17–28.
- Nunes-Nesi, A., Fernie, A.R., Stitt, M., 2010. Metabolic and signaling aspects underpinning the regulation of plant carbon nitrogen interactions. *Mol. Plant* 3, 973–996.
- Oaks, A., 1994. Primary nitrogen assimilation in higher plants and its regulation. *Can. J. Bot. Rev.* 72, 739–750.
- Oberbauer, S.F., Sionit, N., Hastings, S.J., Oechel, W.C., 1986. Effects of CO₂ enrichment and nutrition on growth, photosynthesis, and nutrient concentration of Alaskan tundra plant species. *Can. J. Bot. Rev.* 64, 2993–2998.
- Okada, M., Lieffering, M., Nakamura, H., Yoshimoto, M., Kim, H.Y., Kobayashi, K., 2001. Free-air CO₂ enrichment (FACE) using pure CO₂ injection: system description. *New Phytol.* 150, 251–260.
- Olsson, P., Linder, S., Giesler, R., Hogberg, P., 2005. Fertilization of boreal forest reduces both autotrophic and heterotrophic soil respiration. *Glob. Change Biol.* 11, 1745–1753.
- Ort, D.R., Merchant, S.S., Alric, J., Barkan, A., Blankenship, R.E., Bock, R., Croce, R., Hanson, M.R., Hibberd, J.M., Long, S.P., Moore, T.A., Moroney, J., Niyogi, K.K., Parry, M.A.J., Peralta-Yahya, P.P., Prince, R.C., Redding, K.E., Spalding, M.H., van Wijk, K.J., Vermaas, W.F.J., von Caemmerer, S., Weber, A.P.M., Yeates, T.O., Yuan, J.S., Zhu, X.G., 2015. Redesigning photosynthesis to sustainably meet global food and bioenergy demand. *Proc. Natl. Acad. Sci. U. S. A.* 112, 8529–8536.
- Pachauri, R.K., Allen, M.R., Barros, V.R., Broome, J., Cramer, W., Christ, R., Church, J.A., Clarke, L., Dahe, Q., Dasgupta, P., Dubash, N.K., Edenhofer, O., Elgizouli, I., Field, C.B., Forster, P., Friedlingstein, P., Fuglestad, J., Gomez-Echeverri, L., Hallegatte, S., Hegerl, G., Howden, M., Jiang, K., Jimenez Cisneros, B., Kattsov, V., Lee, H., Mach, K.J., Marotzke, J., Mastrandrea, M.D., Meyer, L., Minx, J., Mulugetta, Y., O'Brien, K., Oppenheimer, M., Pereira, J.J., Pichs-Madruga, R., Plattner, G.-K., Pörtner, H.-O., Power, S.B., Preston, B., Ravindranath, N.H., Reisinger, A., Riahi, K., Rusticucci, N., Scholes, R., Seyboth, K., Sokona, Y., Stavins, R., Stocker, T.F., Tschakert, P., van Vuuren, D., van Ypersele, J.-P., Pachauri, R.K., Meyer, L., 2014. Climate change 2014: synthesis report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change: IPCC.
- Pagani, M., Caldeira, K., Berner, R., Beerling, D.J., 2009. The role of terrestrial plants in limiting atmospheric CO₂ decline over the past 24 million years. *Nature* 460, 85–88.
- Pan, Y., Birdsey, R.A., Fang, J., Houghton, R., Kauppi, P.E., Kurz, W.A., Phillips, O.L., Shvidenko, A., Lewis, S.L., Canadell, J.G., Ciais, P., Jackson, R.B., Pacala, S.W., McGuire, A.D., Piao, S., Rautiainen, A., Sitch, S., Hayes, D., 2011. A large and persistent carbon sink in the world's forests. *Science* 333, 988–993.
- Parton, W.J., Schimel, D.S., Cole, C.V., Ojima, D.S., 1987. Analysis of factors controlling soil organic matter levels in great plains grasslands. *Soil Sci. Soc. Am. J.* 51, 1173–1179.
- Pearson, H.L., Vitousek, P.M., 2001. Stand dynamics, nitrogen accumulation, and symbiotic nitrogen fixation in regenerating stands of *Acacia koa*. *Ecol. Appl.* 11, 1381–1394.
- Phillips, R.P., Bernhardt, E.S., Schlesinger, W.H., 2009. Elevated CO₂ increases root exudation from loblolly pine (*Pinus taeda*) seedlings as an N-mediated response. *Tree Physiol.* 29, 1513–1523.
- Phillips, R.P., Finzi, A.C., Bernhardt, E.S., 2011. Enhanced root exudation induces microbial feedbacks to N cycling in a pine forest under long-term CO₂ fumigation. *Ecol. Lett.* 14, 187–194.
- Pilbeam, D.J., 2015. Breeding crops for improved mineral nutrition under climate change conditions. *J. Exp. Bot.* 66, 3511–3521.
- Pleijel, H., Högy, P., 2015. CO₂ dose-response functions for wheat grain, protein and mineral yield based on FACE and open-top chamber experiments. *Environ. Pollut.* 198, 70–77.
- Pleijel, H., Uddling, J., 2012. Yield vs. quality trade-offs for wheat in response to carbon dioxide and ozone. *Glob. Change Biol.* 18, 596–605.
- Poorter, H., Navas, M.L., 2003. Plant growth and competition at elevated CO₂: on winners, losers and functional groups. *New Phytol.* 157, 175–198.
- Poorter, H., Van Berkel, Y., Baxter, R., Den Hertog, J., Dijkstra, P., Gifford, R.M., Griffin, K.L., Roumet, C., Roy, J., Wong, S.C., 1997. The effect of elevated CO₂ on the chemical composition and construction costs of leaves of 27 C₃ species. *Plant Cell Environ.* 20, 472–482.
- Pritchard, S.G., Strand, A.E., McCormack, M.L., Davis, M.A., Oren, R., 2008. Mycorrhizal and rhizomorph dynamics in a loblolly pine forest during 5 years of free-air-CO₂-enrichment. *Glob. Change Biol.* 14, 1252–1264.
- Rütting, T., Andresen, L.C., 2015. Nitrogen cycle responses to elevated CO₂ depend on ecosystem nutrient status. *Nutr. Cycl. Agroecosyst.* 101, 285–294.
- Raab, T.K., Terry, N., 1994. Nitrogen-source regulation of growth and photosynthesis in *Beta vulgaris* L. *Plant Physiol.* 105, 1159–1166.
- Raich, J.W., Potter, C.S., Bhagawati, D., 2002. Interannual variability in global soil respiration, 1980–94. *Glob. Change Biol.* 8, 800–812.
- Raven, J.A., Wollenweber, B., Handley, L.L., 1992. Ammonia and ammonium fluxes between photolithotrophs and the environment in relation to the global nitrogen cycle. *New Phytol.* 121, 5–18.
- Reay, D.S., Dentener, F., Smith, P., Grace, J., Feely, R.A., 2008. Global nitrogen deposition and carbon sinks. *Nat. Geosci.* 1, 430–437.
- Reich, P.B., 2009. Elevated CO₂ reduces losses of plant diversity caused by nitrogen deposition. *Science* 326, 1399–1402.
- Reich, P.B., Hobbie, S.E., 2013. Decade-long soil nitrogen constraint on the CO₂ fertilization of plant biomass. *Nat. Clim. Change* 3, 278–282.
- Reich, P.B., Oleksyn, J., 2004. Global patterns of plant leaf N and P in relation to temperature and latitude. *Proc. Natl. Acad. Sci. U. S. A.* 101, 11001–11006.
- Reich, P.B., Knops, J., Tilman, D., Craine, J., Ellsworth, D., Tjoelker, M., Lee, T., Wedin, D., Naem, S., Bahaeddin, D., Hendrey, G., Jose, S., Wrage, K., Goth, J., Bengtson, W., 2001. Plant diversity enhances ecosystem responses to elevated CO₂ and nitrogen deposition. *Nature* 410, 809–810.
- Reich, P.B., Hobbie, S.E., Lee, T., Ellsworth, D.S., West, J.B., Tilman, D., Knops, J.M.H., Naeem, S., Trost, J., 2006a. Nitrogen limitation constrains sustainability of ecosystem response to CO₂. *Nature* 440, 922–925.
- Reich, P.B., Hungate, B.A., Luo, Y., 2006b. Carbon-nitrogen interactions in terrestrial ecosystems in response to rising atmospheric carbon dioxide. *Ann. Rev. Ecol. Syst.* 37, 611–636.
- Reich, P.B., Hobbie, S.E., Lee, T.D., 2014. Plant growth enhancement by elevated CO₂ eliminated by joint water and nitrogen limitation. *Nat. Geosci.* 7, 920–924.
- Rennenberg, H., Wildhagen, H., Ehlting, B., 2010. Nitrogen nutrition of poplar trees. *Plant Biol.* 12, 275–291.
- Rice, E.L., Panchoy, S.K., 1972. Inhibition of nitrification by climax ecosystems. *Am. J. Bot.* 59, 1033–1040.
- Richardson, A.E., Lynch, J.P., Ryan, P.R., Delhaize, E., Smith, F.A., Smith, S.E., Harvey, P.R., Ryan, M.H., Veneklaas, E.J., Lambers, H., Oberson, A., 2011. Plant and microbial strategies to improve the phosphorus efficiency of agriculture. *Plant Soil* 349, 121–156.
- Rogers, A., Fischer, B.U., Bryant, J., Frehner, M., Blum, H., Raines, C.A., Long, S.P., 1998. Acclimation of photosynthesis to elevated CO₂ under low-nitrogen nutrition is affected by the capacity for assimilate utilization. Perennial ryegrass under free-air CO₂ enrichment. *Plant Physiol.* 118, 683–689.
- Rogers, A., Gibon, Y., Stitt, M., Morgan, P.B., Bernacchi, C.J., Ort, D.R., Long, S.P., 2006. Increased C availability at elevated carbon dioxide concentration improves N assimilation in a legume. *Plant Cell Environ.* 29, 1651–1658.
- Rogers, A., Ainsworth, E.A., Leakey, A.D.B., 2009. Will elevated carbon dioxide concentration amplify the benefits of nitrogen fixation in legumes? *Plant Physiol.* 151, 1009–1016.
- Rosenthal, D.M., Slattery, R.A., Miller, R.E., Grennan, A.K., Cavagnaro, T.R., Fauquet, C.M., Gleadow, R.M., Ort, D.R., 2012. Cassava about-FACE: greater than expected yield stimulation of cassava (*Manihot esculenta*) by future CO₂ levels. *Glob. Change Biol.* 18, 2661–2675.

- Rubio-Asensio, J.S., Rachmilevitch, S., Bloom, A.J., 2015. Responses of *Arabidopsis* and wheat to rising CO₂ depend on nitrogen source and nighttime CO₂ levels. *Plant Physiol.* 168, 156–163.
- Sage, R.F., 1994. Acclimation of photosynthesis to increasing atmospheric CO₂—the gas-exchange perspective. *Photosynth. Res.* 39, 351–368.
- Sage, R.F., Percy, R.W., Seemann, J.R., 1987. The nitrogen use efficiency of C₃ and C₄ plants. 3. Leaf nitrogen effects on the activity of carboxylating enzymes in *Chenopodium album* L. and *Amaranthus retroflexus* L. *Plant Physiol.* 85, 355–359.
- Sakai, H., Hasegawa, T., Kobayashi, K., 2006. Enhancement of rice canopy carbon gain by elevated CO₂ is sensitive to growth stage and leaf nitrogen concentration. *New Phytol.* 170, 321–332.
- Salsac, L., Chaillou, S., Morotgaudry, J.F., Lesaint, C., 1987. Nitrate and ammonium nutrition in plants. *Plant Physiol. Biochem.* 25, 805–812.
- Sardans, J., Peñuelas, J., 2015. Potassium: a neglected nutrient in global change. *Glob. Ecol. Biogeogr.* 24, 261–275.
- Satapathy, S.S., Swain, D.K., Herath, S., 2014. Field experiments and simulation to evaluate rice cultivar adaptation to elevated carbon dioxide and temperature in sub-tropical India. *Eur. J. Agron.* 54, 21–33.
- Schlesinger, W.H., Andrews, J.A., 2000. Soil respiration and the global carbon cycle. *Biogeochemistry* 48, 7–20.
- Schlesinger, W.H., 2000. Carbon sequestration in soils: some cautions amidst optimism. *Agric. Ecosyst. Environ.* 82, 121–127.
- Schneider, M.K., Luscher, A., Richter, M., Aeschlimann, U., Hartwig, U.A., Blum, H., Frossard, E., Nosberger, J., 2004. Ten years of free-air CO₂ enrichment altered the mobilization of N from soil in *Lolium perenne* L. swards. *Glob. Change Biol.* 10, 1377–1388.
- Schweizer, P., Erisman, K.H., 1985. Effect of nitrate and ammonium nutrition of nonnodulated *Phaseolus vulgaris* L. on phosphoenolpyruvate carboxylase and pyruvate kinase activity. *Plant Physiol.* 78, 455–458.
- Sechley, K.A., Yamaya, T., Oaks, A., 1992. Compartmentation of nitrogen assimilation in higher plants. *Int. Rev. Cytol.* 134, 85–163.
- Shaw, M.R., Zavaleta, E.S., Chiariello, N.R., Cleland, E.E., Mooney, H.A., Field, C.B., 2002. Grassland responses to global environmental changes suppressed by elevated CO₂. *Science* 298, 1987–1990.
- Shimono, H., Okada, M., Yamakawa, Y., Nakamura, H., Kobayashi, K., Hasegawa, T., 2008. Rice yield enhancement by elevated CO₂ is reduced in cool weather. *Glob. Change Biol.* 14, 276–284.
- Sitch, S., Cox, P.M., Collins, W.J., Huntingford, C., 2007. Indirect radiative forcing of climate change through ozone effects on the land-carbon sink. *Nature* 448, 791–794.
- Sitch, S., Friedlingstein, P., Gruber, N., Jones, S., Murray-Tortarolo, G., Ahlström, A., Doney, S.C., Graven, H., Heinze, C., Huntingford, C., 2013. Trends and drivers of regional sources and sinks of carbon dioxide over the past two decades. *Biogeosciences*, 20113–20177 (Discussions 10).
- Smil, V., 2004. Enriching the Earth: Fritz Haber, Carl Bosch, and the Transformation of World Food Production. MIT Press, Cambridge.
- Smith, B.E., 2002. Nitrogenase reveals its inner secrets. *Science* 297, 1654–1655.
- Smith, W.H., Bormann, F.H., Likens, G.E., 1968. Response of chemoautotrophic nitrifiers to forest cutting. *Soil Sci.* 106, 471–473.
- Sokolov, A.P., Kicklighter, D.W., Melillo, J.M., Felzer, B.S., Schlosser, C.A., Cronin, T.W., 2008. Consequences of considering carbon–nitrogen interactions on the feedbacks between climate and the terrestrial carbon cycle. *J. Clim.* 21, 3776–3796.
- Spinnler, D., Egli, P., Körner, C., 2002. Four-year growth dynamics of beech–spruce model ecosystems under CO₂ enrichment on two different forest soils. *Trees* 16, 423–436.
- Spreitzer, R.J., Salvucci, M.E., 2002. Rubisco: structure, regulatory interactions, and possibilities for a better enzyme. *Annu. Rev. Plant Biol.* 53, 449–475.
- Stitt, M., Krapp, A., 1999. The interaction between elevated carbon dioxide and nitrogen nutrition: the physiological and molecular background. *Plant Cell Environ.* 22, 583–621.
- Taub, D.R., Wang, X., 2008. Why are nitrogen concentrations in plant tissues lower under elevated CO₂? A critical examination of the hypotheses. *J. Integr. Plant Biol.* 50, 1365–1374.
- Taub, D.R., Miller, B., Allen, H., 2008. Effects of elevated CO₂ on the protein concentration of food crops: a meta-analysis. *Glob. Change Biol.* 14, 565–575.
- Thornton, P.E., Doney, S.C., Lindsay, K., Moore, J.K., Mahowald, N., Randerson, J.T., Fung, I., Lamarque, J.F., Feddes, J.J., Lee, Y.H., 2009. Carbon–nitrogen interactions regulate climate–carbon cycle feedbacks: results from an atmosphere–ocean general circulation model. *Biogeosciences* 6, 2099–2120.
- Tilman, D., Reich, P.B., Isbell, F., 2012. Biodiversity impacts ecosystem productivity as much as resources, disturbance, or herbivory. *Proc. Natl. Acad. Sci. U. S. A.* 109, 10394–10397.
- Tilman, D., Isbell, F., Cowles, J.M., 2014. Biodiversity and ecosystem functioning. *Annu. Rev. Ecol. Syst.* 45, 471–493.
- Torbert, H.A., Prior, S.A., Rogers, H.H., Runion, G.B., 2004. Elevated atmospheric CO₂ effects on N fertilization in grain sorghum and soybean. *Field Crops Res.* 88, 57–67.
- Tubiello, F.N., Amthor, J.S., Boote, K.J., Donatelli, M., Easterling, W., Fischer, G., Gifford, R.M., Howden, M., Reilly, J., Rosenzweig, C., 2007. Crop response to elevated CO₂ and world food supply—a comment on food for thought... by Long et al. *Science* 312: 1918–1921, 2006. *Eur. J. Agron.* 26, 215–223.
- van Groenigen, K.J., Six, J., Hungate, B.A., de Graaff, M.A., van Breemen, N., van Kessel, C., 2006. Element interactions limit soil carbon storage. *Proc. Natl. Acad. Sci. U. S. A.* 103, 6571–6574.
- Vicca, S., Luysaert, S., Penuelas, J., Campioli, M., Chapin, F.S., Ciais, P., Heinemeyer, A., Hogberg, P., Kutsch, W.L., Law, B.E., Malhi, Y., Papale, D., Piao, S.L., Reichstein, M., Schulze, E.D., Janssens, I.A., 2012. Fertile forests produce biomass more efficiently. *Ecol. Lett.* 15, 520–526.
- Viktor, A., Cramer, M.D., 2005. The influence of root assimilated inorganic carbon on nitrogen acquisition/assimilation and carbon partitioning. *New Phytol.* 165, 157–169.
- Vitousek, P.M., 1984. Litterfall, nutrient cycling, and nutrient limitation in tropical forests. *Ecology* 65, 285–298.
- Vitousek, P.M., Howarth, R.W., 1991. Nitrogen limitation on land and in the sea—how can it occur? *Biogeochemistry* 13, 87–115.
- Vitousek, P.M., Gosz, J.R., Grier, C.C., Melillo, J.M., Reiners, W.A., 1982. A comparative analysis of potential nitrification and nitrate mobility in forest ecosystems. *Ecol. Monogr.* 52, 155–177.
- Vitousek, P.M., Aber, J.D., Howarth, R.W., Likens, G.E., Matson, P.A., Schindler, D.W., Schlesinger, W.H., Tilman, D., 1997. Human alteration of the global nitrogen cycle: sources and consequences. *Ecol. Appl.* 7, 737–750.
- Vitousek, P.M., Cassman, K., Cleveland, C., Crews, T., Field, C.B., Grimm, N.B., Howarth, R.W., Marino, R., Martinelli, L., Rastetter, E.B., Sprent, J.I., 2002. Towards an ecological understanding of biological nitrogen fixation. *Biogeochem* 57, 1–45.
- Wand, S.J.E., Midgley, G.F., Jones, M.H., Curtis, P.S., 1999. Responses of wild C₄ and C₃ grass (*Poaceae*) species to elevated atmospheric CO₂ concentration: a meta-analytic test of current theories and perceptions. *Glob. Change Biol.* 5, 723–741.
- Wang, L., Feng, Z., Schjoerring, J.K., 2013. Effects of elevated atmospheric CO₂ on physiology and yield of wheat (*Triticum aestivum* L.): a meta-analytic test of current hypotheses. *Agric. Ecosyst. Environ.* 178, 57–63.
- Weigel, H.-J., Manderscheid, R., 2012. Crop growth responses to free air CO₂ enrichment and nitrogen fertilization: rotating barley, ryegrass, sugar beet and wheat. *Eur. J. Agron.* 43, 97–107.
- West, J.B., HilleRisLambers, J., Lee, T.D., Hobbie, S.E., Reich, P.B., 2005. Legume species identity and soil nitrogen supply determine symbiotic nitrogen-fixation responses to elevated atmospheric [CO₂]. *New Phytol.* 167, 523–530.
- White, A.C., Rogers, A., Rees, M., Osborne, C.P., 2016. How can we make plants grow faster? A source-sink perspective on growth rate. *J. Exp. Bot.* 67, 31–45.
- Wieder, W.R., Cleveland, C.C., Smith, W.K., Todd-Brown, K., 2015. Future productivity and carbon storage limited by terrestrial nutrient availability. *Nat. Geosci.* 8, 441–444.
- Wong, S.C., 1979. Elevated atmospheric partial pressure of CO₂ and plant growth. 1. Interactions of nitrogen nutrition and photosynthetic capacity in C₃ and C₄ plants. *Oecologia* 44, 68–74.
- Yamakawa, Y., Saigusa, M., Okada, M., Kobayashi, K., 2004. Nutrient uptake by rice and soil solution composition under atmospheric CO₂ enrichment. *Plant Soil* 259, 367–372.
- Yang, L., Liu, H., Wang, Y., Zhu, J., Huang, J., Liu, G., Dong, G., Wang, Y., 2009. Impact of elevated CO₂ concentration on inter-subspecific hybrid rice cultivar Liangyoupeijiu under fully open-air field conditions. *Field Crops Res.* 112, 7–15.
- Yin, X.W., 2002. Responses of leaf nitrogen concentration and specific leaf area to atmospheric CO₂ enrichment: a retrospective synthesis across 62 species. *Glob. Change Biol.* 8, 631–642.
- Yin, X., 2013. Improving ecophysiological simulation models to predict the impact of elevated atmospheric CO₂ concentration on crop productivity. *Ann. Bot.* 112, 465–475.
- Yue, K., Peng, Y., Peng, C., Yang, W., Peng, X., Wu, F., 2016. Stimulation of terrestrial ecosystem carbon storage by nitrogen addition: a meta-analysis. *Sci. Rep.* 6, 19895.
- Zaehle, S., 2013. Terrestrial nitrogen–carbon cycle interactions at the global scale. *Philos. Trans. R. Soc. B* 368.
- Zaehle, S., Friedlingstein, P., Friend, A.D., 2010. Terrestrial nitrogen feedbacks may accelerate future climate change. *Geophys. Res. Lett.* 37.
- Zaehle, S., Jones, C.D., Houlton, B., Lamarque, J.-F., Robertson, E., 2015. Nitrogen availability reduces CMIP5 projections of twenty-first-century land carbon uptake. *J. Clim.* 28, 2494–2511.
- Zak, D.R., Pregitzer, K.S., King, J.S., Holmes, W.E., 2000. Elevated atmospheric CO₂, fine roots and the response of soil microorganisms: a review and hypothesis. *New Phytol.* 147, 201–222.
- Zak, D.R., Holmes, W.E., Pregitzer, K.S., 2007. Atmospheric CO₂ and O₃ alter the flow of N-15 in developing forest ecosystems. *Ecology* 88, 2630–2639.
- Zak, D.R., Pregitzer, K.S., Kubiske, M.E., Burton, A.J., 2011. Forest productivity under elevated CO₂ and O₃: positive feedbacks to soil N cycling sustain decade-long net primary productivity enhancement by CO₂. *Ecol. Lett.* 14, 1220–1226.
- Zanetti, S., Hartwig, U.A., Luscher, A., Hebeisen, T., Frehner, M., Fischer, B.U., Hendrey, G.R., Blum, H., Nosberger, J., 1996. Stimulation of symbiotic N₂ fixation in *Trifolium repens* L. under elevated atmospheric pCO₂ in a grassland ecosystem. *Plant Physiol.* 112, 575–583.
- Zavaleta, E.S., Shaw, M.R., Chiariello, N.R., Mooney, H.A., Field, C.B., 2003. Additive effects of simulated climate changes, elevated CO₂, and nitrogen deposition on grassland diversity. *Proc. Natl. Acad. Sci. U. S. A.* 100, 7650–7654.
- Ziska, L.H., Bunce, J.A., 2007. Predicting the impact of changing CO₂ on crop yields: some thoughts on food. *New Phytol.* 175, 607–617.